

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
11 November 2004 (11.11.2004)

PCT

(10) International Publication Number  
**WO 2004/096767 A1**

(51) International Patent Classification<sup>7</sup>: **C07D 209/08**,  
409/06, 401/06, 403/06, 407/06, 417/06, A61K 31/404,  
A61P 25/08

**William, Patrick** [GB/DK]; Jyllingevej 81 st, DK-2720  
Vanløse (DK).

(21) International Application Number:  
PCT/DK2004/000283

(22) International Filing Date: 23 April 2004 (23.04.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
PA200300631 25 April 2003 (25.04.2003) DK  
60/465,387 25 April 2003 (25.04.2003) US

(71) Applicant (for all designated States except US): **H. LUNDBECK A/S** [DK/DK]; Ottiliavej 9, DK-2500  
Valby-Copenhagen (DK).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **KHANZHIN, Nikolay** [RU/DK]; Finsensvej 44A, 4. tv., DK-2000  
Frederiksberg (DK). **ROTTLANDER, Mario** [DE/DK];  
Svanemosen 10, DK-2670 Greve (DK). **WATSON,**

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: SUBSTITUTED INDOLINE AND INDOLE DERIVATIVES



(57) Abstract: The present invention relates to aniline derivatives of the general formula I or pharmaceutically acceptable salts thereof, [Formula (I)], and their use.

WO 2004/096767 A1

## Substituted indoline and indole derivatives

### Field of the invention

The present invention relates to novel substituted indole and indoline derivatives  
5 being openers of the KCNQ family potassium ion channels. The compounds are  
useful for the prevention, treatment and inhibition of disorders and diseases being  
responsive to opening of the KCNQ family potassium ion channels, one such disease  
is epilepsy.

### 10 Background of the invention

Ion channels are cellular proteins that regulate the flow of ions, including potassium,  
calcium, chloride and sodium into and out of cells. Such channels are present in all  
animal and human cells and affect a variety of processes including neuronal  
transmission, muscle contraction, and cellular secretion.

15 Humans have over 70 genes encoding potassium channel subtypes (Jentsch *Nature  
Reviews Neuroscience* 2000, 1, 21-30) with a great diversity with regard to both  
structure and function. Neuronal potassium channels, which are found in the brain, are  
primarily responsible for maintaining a negative resting membrane potential, as well  
20 as controlling membrane repolarisation following an action potential.

One subset of potassium channel genes is the KCNQ family. Mutations in four out of  
five KCNQ genes have been shown to underlie diseases including cardiac  
arrhythmias, deafness and epilepsy (Jentsch *Nature Reviews Neuroscience* 2000, 1,  
25 21-30).

The KCNQ4 gene is thought to encode the molecular correlate of a potassium channel  
found in outer hair cells of the cochlea and in Type I hair cells of the vestibular  
apparatus, in which, mutations can lead to a form of inherited deafness.

30 KCNQ1 (KvLQT1) is co-assembled with the product of the KCNE1 (minimal K(+)-  
channel protein) gene in the heart to form a cardiac-delayed rectifier-like K(+)-  
current. Mutations in this channel can cause one form of inherited long QT syndrome

type 1 (LQT1), as well as being associated with a form of deafness (Robbins *Pharmacol Ther* 2001, 90, 1-19).

5 The genes KCNQ2 and KCNQ3 were discovered in 1988 and appear to be mutated in an inherited form of epilepsy known as benign familial neonatal convulsions (Rogawski *Trends in Neurosciences* 2000, 23, 393-398). The proteins encoded by the KCNQ2 and KCNQ3 genes are localised in the pyramidal neurons of the human cortex and hippocampus, regions of the brain associated with seizure generation and propagation (Cooper et al. *Proceedings National Academy of Science U S A* 2000, 97, 10 4914-4919).

KCNQ2 and KCNQ3 are two potassium channel subunits that form "M-currents" when expressed in vitro. The M-current is a non-inactivating potassium current found in many neuronal cell types. In each cell type, it is dominant in controlling membrane 15 excitability by being the only sustained current in the range of action potential initiation (Marrion *Annual Review Physiology* 1997, 59, 483-504). Modulation of the M-current has dramatic effects on neuronal excitability, for example activation of the current will reduce neuronal excitability. Openers of these KCNQ channels, or activators of the M-current, will reduce excessive neuronal activity and may thus be 20 of use in the treatment, prevention or inhibition of seizures and other diseases and disorders characterised by excessive neuronal activity, such as neuronal hyperexcitability including convulsive disorders, epilepsy and neuropathic pain.

Retigabine (D-23129; N-(2-amino-4-(4-fluorobenzylamino)-phenyl) carbamic acid 25 ethyl ester) and analogues thereof are disclosed in EP554543. Retigabine is an anti-convulsive compound with a broad spectrum and potent anticonvulsant properties, both in vitro and in vivo. It is active after oral and intraperitoneal administration in rats and mice in a range of anticonvulsant tests including: electrically induced seizures, seizures induced chemically by pentylenetetrazole, picrotoxin and N-methyl- 30 D-aspartate (NMDA) and in a genetic animal model, the DBA/2 mouse (Rostock et al. *Epilepsy Research* 1996, 23, 211-223). In addition, retigabine is active in the amygdala kindling model of complex partial seizures, further indicating that this compound has potential for anti-convulsive therapy. In clinical trials, retigabine has

recently shown effectiveness in reducing the incidence of seizures in epileptic patients (Bialer et al. *Epilepsy Research* **2002**, 51, 31-71).

Retigabine has been shown to activate a K(+) current in neuronal cells and the  
5 pharmacology of this induced current displays concordance with the published  
pharmacology of the M-channel, which recently was correlated to the KCNQ2/3 K(+) channel heteromultimer. This suggests that activation of KCNQ2/3 channels may be responsible for some of the anticonvulsant activity of this agent (Wickenden et al. *Molecular Pharmacology* **2000**, 58, 591-600) – and that other agents working by the  
10 same mechanism may have similar uses.

KCNQ 2 and 3 channels have also been reported to be upregulated in models of neuropathic pain (Wickenden et al. *Society for Neuroscience Abstracts* **2002**, 454.7), and potassium channel modulators have been hypothesised to be active in both  
15 neuropathic pain and epilepsy (Schroder et al. *Neuropharmacology* **2001**, 40, 888-898).

Retigabine has also been shown to be beneficial in animal models of neuropathic pain (Blackburn-Munro and Jensen *European Journal of Pharmacology* **2003**, 460, 109-  
20 116), and it is thus suggested that openers of KCNQ channels will be of use in treating pain disorders including neuropathic pain.

The localisation of KCNQ channel mRNA is reported in brain and other central nervous system areas associated with pain (Goldstein et al. *Society for Neuroscience Abstracts* **2003**, 53.8).  
25

In addition to a role in neuropathic pain, the expression of mRNA for KCNQ 2-5 in the trigeminal and dorsal root ganglia and in the trigeminal nucleus caudalis implies that openers of these channels may also affect the sensory processing of migraine pain  
30 (Goldstein et al. *Society for Neuroscience Abstracts* **2003**, 53.8).

Recent reports demonstrate that mRNA for KCNQ 3 and 5, in addition to that for KCNQ2, are expressed in astrocytes and glial cells. Thus KCNQ 2, 3 and 5 channels

may help modulate synaptic activity in the CNS and contribute to the neuroprotective effects of KCNQ channel openers (Noda et al., *Society for Neuroscience Abstracts* 2003, 53.9).

5 Retigabine and other KCNQ modulators may thus exhibit protection against the neurodegenerative aspects of epilepsy, as retigabine has been shown to prevent limbic neurodegeneration and the expression of markers of apoptosis following kainic acid-induced status epilepticus in the rat (Ebert et al. *Epilepsia* 2002, 43 Suppl 5, 86-95). This may have relevance for preventing the progression of epilepsy in patients, i.e. be  
10 anti-epileptogenic. Retigabine has also been shown to delay the progression of hippocampal kindling in the rat, a further model of epilepsy development (Tober et al. *European Journal Of Pharmacology* 1996, 303, 163-169).

It is thus suggested that these properties of retigabine and other KCNQ modulators  
15 may prevent neuronal damage induced by excessive neuronal activation, and may be of use in the treatment of neurodegenerative diseases, and be disease modifying (or antiepileptogenic) in patients with epilepsy.

Given that anticonvulsant compounds such as benzodiazepines and chlormethiazole  
20 are used clinically in the treatment of the ethanol withdrawal syndrome and that other anticonvulsant compounds e.g. gabapentin, are very effective in animal models of this syndrome (Watson et al. *Neuropharmacology* 1997, 36, 1369-1375), other anticonvulsant compounds such as KCNQ openers are thus expected to be effective in this condition.

25 mRNA for KCNQ 2 and 3 subunits are found in brain regions associated with anxiety and emotional behaviours such as bipolar disorder e.g. hippocampus and amygdala (Saganich et al. *Journal of Neuroscience* 2001, 21, 4609-4624), and retigabine is reportedly active in some animal models of anxiety-like behaviour (Hartz et al.  
30 *Journal of Psychopharmacology* 2003, 17 suppl 3, A28,B16), and other clinically used anticonvulsant compounds are used in the treatment of bipolar disorder.

WO 200196540 discloses the use of modulators of the M-current formed by expression of KCNQ2 and KCNQ3 genes for insomnia, while WO 2001092526 discloses that modulators of KCNQ5 can be utilized for the treatment of sleep disorders.

5

WO01/022953 describes the use of retigabine for prophylaxis and treatment of neuropathic pain such as allodynia, hyperalgesic pain, phantom pain, neuropathic pain related to diabetic neuropathy and neuropathic pain related to migraine.

- 10 WO02/049628 describes the use of retigabine for the prevention, treatment, inhibition and amelioration of anxiety disorders such as anxiety, generalized anxiety disorder, panic anxiety, obsessive compulsive disorder, social phobia, performance anxiety, post-traumatic stress disorder, acute stress reaction, adjustment disorders, hypochondriacal disorders, separation anxiety disorder, agoraphobia and specific  
15 phobias.

- WO97/15300 describes the use of retigabine for the treatment of neurodegenerative disorders such as Alzheimer's disease; Huntington's chorea; sclerosis such as multiple sclerosis and amyotrophic lateral sclerosis; Creutzfeld-Jakob disease; Parkinson's  
20 disease; encephalopathies induced by AIDS or infection by rubella viruses, herpes viruses, borrelia and unknown pathogens; trauma-induced neurodegenerations; neuronal hyperexcitation states such as in medicament withdrawal or intoxication; and neurodegenerative diseases of the peripheral nervous system such as polyneuropathies and polyneuritides.

25

Hence, there is a great desire for novel compounds, which are potent openers of the KCNQ family potassium channels.

- Also desired are novel compounds with improved properties relative to known  
30 compounds, which are openers of the KCNQ family potassium channels, such as retigabine. Improvement of one or more of the following parameters is desired: half-life, clearance, selectivity, interactions with other medications, bioavailability, potency, formulability, chemical stability, metabolic stability, membrane

permeability, solubility and therapeutic index. The improvement of such parameters may lead to improvements such as:


- an improved dosing regime by reducing the number of required doses a day,
- ease of administration to patients on multiple medications,
- 5   • reduced side effects,
- enlarged therapeutic index,
- improved tolerability or
- improved compliance.

10

### Summary of the invention

One object of the present invention is to provide novel compounds, which are potent openers of the KCNQ family potassium channels.

- 15   The compounds of the invention are substituted indoline and indole derivatives of the general formula **I** or salts thereof



20


wherein the dotted line,  $q$ ,  $s$ ,  $U$ ,  $Y$ ,  $X$ ,  $Z$ ,  $R^1$ ,  $R^{1'}$ ,  $R^2$  and  $R^3$  are as defined below.

25

The invention further relates to a pharmaceutical composition comprising a compound of formula I, and the use thereof.

## 5 Description of the invention

Accordingly, the present invention relates to substituted indole and indoline derivatives of the general formula I



10 wherein

the dotted line represents an optional bond;

$R^1$  and  $R^{1'}$  are independently selected from the group consisting of hydrogen,  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, hydroxy- $C_{1-6}$ -alk(en/yn)yl, hydroxy- $C_{3-8}$ -cycloalk(en)yl, hydroxy- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, halo- $C_{1-6}$ -alk(en/yn)yl, halo- $C_{3-8}$ -cycloalk(en)yl, halo- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, cyano- $C_{1-6}$ -alk(en/yn)yl, cyano- $C_{3-8}$ -cycloalk(en)yl and cyano- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl; or

20  $R^1$  and  $R^{1'}$  together with the carbon atom to which they are attached form a 3-8 membered saturated or unsaturated ring which optionally contains 1 or 2 heteroatoms;

$s$  is 0 or 1;

25  $U$  is O,  $NR^{11}$ , S,  $SO_2$ ,  $SO_2NR^{11}$ , CO-O or CO- $NR^{11}$ ; wherein  $R^{11}$  is selected from the group consisting of hydrogen,  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl,  $C_{3-8}$ -



cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl; or **R**<sup>2</sup> and **R**<sup>11</sup> together with the nitrogen atom to which they are attached form a 4-8 membered saturated or unsaturated ring which optionally contains 1, 2 or 3 further heteroatoms;

- 5 **R**<sup>2</sup> is selected from the group consisting of hydrogen, C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, Ar, Ar-C<sub>1-6</sub>-alk(en/yn)yl, Ar-C<sub>3-8</sub>-cycloalk(en)yl, Ar-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, acyl, hydroxy-C<sub>1-6</sub>-alk(en/yn)yl, hydroxy-C<sub>3-8</sub>-cycloalk(en)yl, hydroxy-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, halogen, halo-C<sub>1-6</sub>-alk(en/yn)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, cyano, cyano-C<sub>1-6</sub>-alk(en/yn)yl, cyano-C<sub>3-8</sub>-cycloalk(en)yl, cyano-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, -NO<sub>2</sub>, **NR**<sup>10</sup>**R**<sup>10'</sup>-C<sub>1-6</sub>-alk(en/yn)yl, **NR**<sup>10</sup>**R**<sup>10'</sup>-C<sub>3-8</sub>-cycloalk(en)yl and **NR**<sup>10</sup>**R**<sup>10'</sup>-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl; wherein

- 15 **R**<sup>10</sup> and **R**<sup>10'</sup> are independently selected from the group consisting of hydrogen, C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, hydroxy-C<sub>1-6</sub>-alk(en/yn)yl, hydroxy-C<sub>3-8</sub>-cycloalk(en)yl, hydroxy-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, halo-C<sub>1-6</sub>-alk(en/yn)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, cyano-C<sub>1-6</sub>-alk(en/yn)yl, cyano-C<sub>3-8</sub>-cycloalk(en)yl and cyano-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, or

- 20 **R**<sup>10</sup> and **R**<sup>10'</sup> together with the nitrogen atom to which they are attached form a 4-8 membered saturated or unsaturated ring which optionally contains 1, 2 or 3 further heteroatoms;

with the proviso that when **R**<sup>2</sup> is NO<sub>2</sub>, halogen or cyano then s is 0; and

- 25 with the proviso that when **R**<sup>2</sup> is a hydrogen atom or acyl and s is 1 then **U** is **NR**<sup>11</sup>, O or S;

wherein the group -(**U**)<sub>s</sub>-**R**<sup>2</sup> is linked to position 4 or 6 of the indole or indoline;

**q** is 0 or 1;

30

**Z** is O or S;

**X** is CO or SO<sub>2</sub>; with the proviso that **q** is 0 when **X** is SO<sub>2</sub>;

$R^3$  is selected from the group consisting of  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl, heterocycloalk(en)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl,  $C_{1-6}$ -alk(en/yn)yl- $C_{3-8}$ -cycloalk(en)yl,  $C_{1-6}$ -alk(en/yn)yl-heterocycloalk(en)yl, Ar, Ar- $C_{1-6}$ -alk(en/yn)yl, Ar- $C_{3-8}$ -cycloalk(en)yl, Ar-heterocycloalk(en)yl, Ar- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, Ar- $C_{1-6}$ -alk(en/yn)yl- $C_{3-8}$ -cycloalk(en)yl, Ar- $C_{1-6}$ -alk(en/yn)yl-heterocycloalk(en)yl,  $C_{1-6}$ -alk(en/yn)yoxy- $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yoxy- $C_{1-6}$ -alk(en/yn)yl,  $C_{1-6}$ -alk(en/yn)yoxy- $C_{3-8}$ -cycloalk(en)yl,  $C_{1-6}$ -alk(en/yn)yoxy-heterocycloalk(en)yl, Ar-oxy- $C_{1-6}$ -alk(en/yn)yl, Ar- $C_{1-6}$ -alk(en/yn)yoxy- $C_{1-6}$ -alk(en/yn)yl,  $C_{1-6}$ -alk(en/yn)yoxy-carbonyl- $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yoxy-carbonyl- $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yoxy-carbonyl- $C_{1-6}$ -alk(en/yn)yl, hydroxy- $C_{1-6}$ -alk(en/yn)yl, hydroxy- $C_{3-8}$ -cycloalk(en)yl, hydroxy-heterocycloalk(en)yl, hydroxy- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, hydroxy- $C_{1-6}$ -alk(en/yn)yl- $C_{3-8}$ -cycloalk(en)yl, hydroxy- $C_{1-6}$ -alk(en/yn)yl-heterocycloalk(en)yl, halo- $C_{1-6}$ -alk(en/yn)yl, halo- $C_{3-8}$ -cycloalk(en)yl, halo-heterocycloalk(en)yl, halo- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, halo- $C_{1-6}$ -alk(en/yn)yl- $C_{3-8}$ -cycloalk(en)yl, halo- $C_{1-6}$ -alk(en/yn)yl-heterocycloalk(en)yl, halo- $C_{1-6}$ -alk(en/yn)yl-Ar, halo- $C_{3-8}$ -cycloalk(en)yl-Ar, halo- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl-Ar, halo- $C_{1-6}$ -alk(en/yn)yl- $C_{3-8}$ -cycloalk(en)yl-Ar, cyano- $C_{1-6}$ -alk(en/yn)yl, cyano- $C_{3-8}$ -cycloalk(en)yl, cyano-heterocycloalk(en)yl, cyano- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, cyano- $C_{1-6}$ -alk(en/yn)yl- $C_{3-8}$ -cycloalk(en)yl, cyano- $C_{1-6}$ -alk(en/yn)yl-heterocycloalk(en)yl and - $NR^{12}R^{12'}$ , optionally substituted  $NR^{12}R^{12'}$ - $C_{1-6}$ -alk(en/yn)yl, optionally substituted  $NR^{12}R^{12'}$ - $C_{3-8}$ -cycloalk(en)yl, optionally substituted  $NR^{12}R^{12'}$ - $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl; wherein

$R^{12}$  and  $R^{12'}$  are independently selected from the group consisting of hydrogen,  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, Ar, Ar- $C_{1-6}$ -alk(en/yn)yl, Ar- $C_{3-8}$ -cycloalk(en)yl, Ar- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, hydroxy- $C_{1-6}$ -alk(en/yn)yl, hydroxy- $C_{3-8}$ -cycloalk(en)yl, hydroxy- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, halo- $C_{1-6}$ -alk(en/yn)yl, halo- $C_{3-8}$ -cycloalk(en)yl, halo- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, cyano- $C_{1-6}$ -alk(en/yn)yl, cyano- $C_{3-8}$ -cycloalk(en)yl and cyano- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, or


$R^{12}$  and  $R^{12'}$  together with the nitrogen atom to which they are attached form a 4-8 membered saturated or unsaturated ring which optionally contains 1, 2 or 3 further heteroatoms;

with the proviso that when  $R^3$  is  $NR^{12}R^{12'}$  then  $q$  is 0;


5

and


$Y$  represents a group of formula II, III, IV, V, VI, XXX and XXXI:




II




III




IV



V




VI



XXX

or



XXXI

wherein

the line represents a bond attaching the group represented by Y to the carbon atom;

5 W is O or S;

T is N, NH or O;

L is N, C or CH;

10

a is 0, 1, 2 or 3;

b is 0, 1, 2, 3 or 4;

15 c is 0 or 1;

d is 0, 1, 2 or 3;

e is 0, 1 or 2;

20

f is 0, 1, 2, 3, 4 or 5;

g is 0, 1, 2, 3 or 4;

25 h is 0, 1, 2 or 3;

j is 0, 1, 2 or 3; with the proviso that when T is a nitrogen atom then j is 0, 1, 2 or 3;  
and when T is NH or an oxygen atom then j is 0, 1 or 2;

30 k is 0, 1, 2, 3 or 4; and

each R<sup>5</sup> is independently selected from the group consisting of a C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, Ar, Ar-C<sub>1-6</sub>-alk(en/yn)yl, Ar-

thio, Ar-oxy, acyl, C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, halogen, halo-C<sub>1-6</sub>-alk(en/yn)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, -CO-NR<sup>6</sup>R<sup>6'</sup>, cyano, cyano-C<sub>1-6</sub>-alk(en/yn)yl, cyano-C<sub>3-8</sub>-cycloalk(en)yl, cyano-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, -NR<sup>7</sup>R<sup>7'</sup>, -S-  
 5 R<sup>8</sup> and -SO<sub>2</sub>R<sup>8</sup>, or

two adjacent R<sup>5</sup> together with the aromatic group to which they are attached form a 4-8 membered ring which optionally contains one or two heteroatoms;

R<sup>6</sup> and R<sup>6'</sup> are independently selected from the group consisting of hydrogen, C<sub>1-6</sub>-  
 10 alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl and Ar;

R<sup>7</sup> and R<sup>7'</sup> are independently selected from the group consisting of hydrogen, C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, Ar and acyl;


15 and

R<sup>8</sup> is selected from the group consisting of hydrogen, C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, Ar and -NR<sup>9</sup>R<sup>9'</sup>; wherein  
 R<sup>9</sup> and R<sup>9'</sup> are independently selected from the group consisting of hydrogen, C<sub>1-6</sub>-  
 20 alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl and C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl;  
 provided that when R<sup>8</sup> is -NR<sup>9</sup>R<sup>9'</sup> then R<sup>5</sup> is not -S-R<sup>8</sup>;

or salts thereof.

25

A particular embodiment of the invention relates to substituted indole and indoline derivatives of the general formula I



(I)

wherein

the dotted line represents an optional bond;

5

$R^1$  and  $R^{1'}$  are independently selected from the group consisting of hydrogen,  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, hydroxy- $C_{1-6}$ -alk(en/yn)yl, hydroxy- $C_{3-8}$ -cycloalk(en)yl, hydroxy- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, halo- $C_{1-6}$ -alk(en/yn)yl, halo- $C_{3-8}$ -cycloalk(en)yl, halo- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, cyano- $C_{1-6}$ -alk(en/yn)yl, cyano- $C_{3-8}$ -cycloalk(en)yl and cyano- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl; or

10

$R^1$  and  $R^{1'}$  form a 3-8 membered saturated or unsaturated ring which optionally contains 1 or 2 further heteroatoms;

15

s is 0 or 1;

U is O,  $NR^{11}$ , S,  $SO_2$ ,  $SO_2NR^{11}$ , CO-O or CO- $NR^{11}$ ; wherein  $R^{11}$  is selected from the group consisting of hydrogen,  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl; or

20

$R^2$  and  $R^{11}$  together with the nitrogen atom to which they are attached form a 4-8 membered saturated or unsaturated ring which optionally contains 1, 2 or 3 further heteroatoms;

25

$R^2$  is selected from the group consisting of hydrogen,  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, Ar, Ar- $C_{1-6}$ -alk(en/yn)yl, Ar- $C_{3-8}$ -cycloalk(en)yl, Ar- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, acyl, hydroxy- $C_{1-6}$ -

- alk(en/yn)yl, hydroxy-C<sub>3-8</sub>-cycloalk(en)yl, hydroxy-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, halo-C<sub>1-6</sub>-alk(en/yn)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, cyano, cyano-C<sub>1-6</sub>-alk(en/yn)yl, cyano-C<sub>3-8</sub>-cycloalk(en)yl, cyano-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, -NO<sub>2</sub>, NR<sup>10</sup>R<sup>10'</sup>-C<sub>1-6</sub>-alk(en/yn)yl, NR<sup>10</sup>R<sup>10'</sup>-C<sub>3-8</sub>-cycloalk(en)yl and NR<sup>10</sup>R<sup>10'</sup>-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl; wherein
- R<sup>10</sup> and R<sup>10'</sup> are independently selected from the group consisting of hydrogen, C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, hydroxy-C<sub>1-6</sub>-alk(en/yn)yl, hydroxy-C<sub>3-8</sub>-cycloalk(en)yl, hydroxy-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, halo-C<sub>1-6</sub>-alk(en/yn)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, cyano-C<sub>1-6</sub>-alk(en/yn)yl, cyano-C<sub>3-8</sub>-cycloalk(en)yl and cyano-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, or
- R<sup>10</sup> and R<sup>10'</sup> together with the nitrogen atom to which they are attached form a 4-8 membered saturated or unsaturated ring which optionally contains 1, 2 or 3 further heteroatoms;
- with the proviso that when R<sup>2</sup> is NO<sub>2</sub>, halogen or cyano then s is 0; and
- with the proviso that when R<sup>2</sup> is a hydrogen atom or acyl and s is 1 then U is NR<sup>11</sup>, O or S;
- wherein the group -(U)<sub>s</sub>-R<sup>2</sup> is linked to position 4 or 6 of the indole or indoline;
- q is 0 or 1;
- Z is O or S;
- X is CO or SO<sub>2</sub>; with the proviso that q is 0 when X is SO<sub>2</sub>;
- R<sup>3</sup> is selected from the group consisting of C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl, heterocycloalk(en)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl, C<sub>1-6</sub>-alk(en/yn)yl-heterocycloalk(en)yl, Ar, Ar-C<sub>1-6</sub>-alk(en/yn)yl, Ar-C<sub>3-8</sub>-cycloalk(en)yl, Ar-heterocycloalk(en)yl, Ar-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, Ar-C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl, Ar-C<sub>1-6</sub>-alk(en/yn)yl-heterocycloalk(en)yl, C<sub>1-6</sub>-alk(en/yn)yl-oxy-C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl-oxy-

$C_{1-6}$ -alk(en/yn)yl,  $C_{1-6}$ -alk(en/yn)yoxy- $C_{3-8}$ -cycloalk(en)yl,  $C_{1-6}$ -alk(en/yn)yoxy-heterocycloalk(en)yl, Ar-oxy- $C_{1-6}$ -alk(en/yn)yl, Ar- $C_{1-6}$ -alk(en/yn)yoxy- $C_{1-6}$ -alk(en/yn)yl,  $C_{1-6}$ -alk(en/yn)yoxy-carbonyl- $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yoxy-carbonyl- $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yoxy-carbonyl- $C_{1-6}$ -alk(en/yn)yl, hydroxy- $C_{1-6}$ -alk(en/yn)yl, hydroxy- $C_{3-8}$ -cycloalk(en)yl, hydroxy-heterocycloalk(en)yl, hydroxy- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, hydroxy- $C_{1-6}$ -alk(en/yn)yl- $C_{3-8}$ -cycloalk(en)yl, hydroxy- $C_{1-6}$ -alk(en/yn)yl-heterocycloalk(en)yl, halo- $C_{1-6}$ -alk(en/yn)yl, halo- $C_{3-8}$ -cycloalk(en)yl, halo-heterocycloalk(en)yl, halo- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, halo- $C_{1-6}$ -alk(en/yn)yl- $C_{3-8}$ -cycloalk(en)yl, halo- $C_{1-6}$ -alk(en/yn)yl-heterocycloalk(en)yl, halo- $C_{1-6}$ -alk(en/yn)yl-Ar, halo- $C_{3-8}$ -cycloalk(en)yl-Ar, halo- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl-Ar, halo- $C_{1-6}$ -alk(en/yn)yl- $C_{3-8}$ -cycloalk(en)yl-Ar, cyano- $C_{1-6}$ -alk(en/yn)yl, cyano- $C_{3-8}$ -cycloalk(en)yl, cyano-heterocycloalk(en)yl, cyano- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, cyano- $C_{1-6}$ -alk(en/yn)yl- $C_{3-8}$ -cycloalk(en)yl, cyano- $C_{1-6}$ -alk(en/yn)yl-heterocycloalk(en)yl, acyl- $C_{1-6}$ -alk(en/yn)yl, acyl- $C_{3-8}$ -cycloalk(en)yl, acyl-heterocycloalk(en)yl, acyl- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, acyl- $C_{1-6}$ -alk(en/yn)yl- $C_{3-8}$ -cycloalk(en)yl, acyl- $C_{1-6}$ -alk(en/yn)yl-heterocycloalk(en)yl and - $NR^{12}R^{12'}$ ; wherein

$R^{12}$  and  $R^{12'}$  are independently selected from the group consisting of hydrogen,  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, Ar, Ar- $C_{1-6}$ -alk(en/yn)yl, Ar- $C_{3-8}$ -cycloalk(en)yl, Ar- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, hydroxy- $C_{1-6}$ -alk(en/yn)yl, hydroxy- $C_{3-8}$ -cycloalk(en)yl, hydroxy- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, halo- $C_{1-6}$ -alk(en/yn)yl, halo- $C_{3-8}$ -cycloalk(en)yl, halo- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, cyano- $C_{1-6}$ -alk(en/yn)yl, cyano- $C_{3-8}$ -cycloalk(en)yl and cyano- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, or


$R^{12}$  and  $R^{12'}$  together with the nitrogen atom to which they are attached form a 4-8 membered saturated or unsaturated ring which optionally contains 1, 2 or 3 further heteroatoms;

with the proviso that when  $R^3$  is  $NR^{12}R^{12'}$  then q is 0;


and

Y represents a group of formula II, III, IV, V and VI:







II




III



IV



V



VI

wherein

5

the line represents a bond attaching the group represented by Y to the carbon atom;

W is O or S;

10 a is 0, 1, 2 or 3;

b is 0, 1, 2, 3 or 4;

c is 0 or 1;

15

d is 0, 1, 2 or 3;



One embodiment of the invention relates to compounds of formula I, wherein the dotted line represents a bond.

Another embodiment of the invention relates to compounds of formula I, wherein the  
5 dotted line does not represent a bond.

One further embodiment of the invention relates to compounds of formula I, wherein  $R^1$  and  $R^{1'}$  are independently selected from the group consisting of hydroxy- $C_{1-6}$ -  
10 alk(en/yn)yl, hydroxy- $C_{3-8}$ -cycloalk(en)yl, hydroxy- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, halo- $C_{1-6}$ -alk(en/yn)yl, halo- $C_{3-8}$ -cycloalk(en)yl, halo- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, cyano- $C_{1-6}$ -alk(en/yn)yl, cyano- $C_{3-8}$ -cycloalk(en)yl and cyano- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl.

15 Another embodiment of the invention relates to compounds of formula I, wherein  $R^1$  and  $R^{1'}$  are independently selected from the group consisting of hydrogen,  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl and  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl.

A further embodiment of the invention relates to compounds of formula I, wherein  $R^1$   
20 and  $R^{1'}$  form a 3-8 membered saturated or unsaturated ring which optionally contains 1 or 2 heteroatoms. In a further embodiment the 3-8 membered saturated or unsaturated ring is a saturated carbocyclic ring, typically cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl.

25 Yet another embodiment of the invention relates to compounds of formula I, wherein  $R^1$  and  $R^{1'}$  are independently selected from the group consisting of hydrogen and  $C_{1-6}$ -alk(en/yn)yl.

In yet another embodiment, the invention relates to compounds of formula I, wherein  
30 at least one of  $R^1$  and  $R^{1'}$  is  $C_{1-6}$ -alk(en/yn)yl, typically  $C_{1-3}$ -alk(en/yn)yl.

In a preferred embodiment, the invention relates to compounds of formula I, wherein  $R^1$  or  $R^{1'}$  is a hydrogen atom.

In a preferred embodiment, the invention relates to compounds of formula **I**, wherein at least one of **R**<sup>1</sup> and **R**<sup>1'</sup> is a hydrogen atom.

5 In a more preferred embodiment, the invention relates to compounds of formula **I**, wherein both **R**<sup>1</sup> and **R**<sup>1'</sup> are hydrogen atoms.

In a preferred embodiment, the invention relates to compounds of formula **I**, wherein **s** is 0.

10 In another preferred embodiment, the invention relates to compounds of formula **I**, wherein **s** is 1.

In one embodiment, the invention relates to compounds of formula **I**, wherein **s** is 1 and **U** is O.

15 In another embodiment, the invention relates to compounds of formula **I**, wherein **s** is 1 and **U** is S.

20 In yet another embodiment, the invention relates to compounds of formula **I**, wherein **s** is 1 and **U** is SO<sub>2</sub>.

In yet another embodiment, the invention relates to compounds of formula **I**, wherein **s** is 1 and **U** is SO<sub>2</sub>NR<sup>11</sup>.

25 In yet another embodiment, the invention relates to compounds of formula **I**, wherein **s** is 1 and **U** is CO-O.

In yet another embodiment, the invention relates to compounds of formula **I**, wherein **s** is 1 and **U** is CO-NR<sup>11</sup>.

30 In a preferred embodiment, the invention relates to compounds of formula **I**, wherein **s** is 1 and **U** is NR<sup>11</sup>.

In yet another embodiment, the invention relates to compounds of formula I, wherein s is 1 and U is  $\text{SO}_2\text{NR}^{11}$ ,  $\text{CO-NR}^{11}$  or  $\text{NR}^{11}$  and  $\text{R}^{11}$  is a hydrogen atom.

In a preferred embodiment, the invention relates to compounds of formula I, wherein  
5 s is 1 and U is  $\text{NR}^{11}$  and  $\text{R}^{11}$  is a hydrogen atom.

One embodiment of the invention relates to compounds of formula I, wherein  $\text{R}^2$  is selected from the group consisting of acyl, hydroxy- $\text{C}_{1-6}$ -alk(en/yn)yl, hydroxy- $\text{C}_{3-8}$ -cycloalk(en)yl, hydroxy- $\text{C}_{3-8}$ -cycloalk(en)yl- $\text{C}_{1-6}$ -alk(en/yn)yl, cyano- $\text{C}_{1-6}$ -  
10 alk(en/yn)yl, cyano- $\text{C}_{3-8}$ -cycloalk(en)yl, cyano- $\text{C}_{3-8}$ -cycloalk(en)yl- $\text{C}_{1-6}$ -alk(en/yn)yl,  $\text{NR}^{10}\text{R}^{10'}$ - $\text{C}_{1-6}$ -alk(en/yn)yl,  $\text{NR}^{10}\text{R}^{10'}$ - $\text{C}_{3-8}$ -cycloalk(en)yl and  $\text{NR}^{10}\text{R}^{10'}$ - $\text{C}_{3-8}$ -cycloalk(en)yl- $\text{C}_{1-6}$ -alk(en/yn)yl;

with the proviso that when  $\text{R}^2$  is acyl and s is 1 then U is  $\text{NR}^{11}$ , O or S.

15 Another embodiment of the invention relates to compounds of formula I, wherein  $\text{R}^2$  is selected from the group consisting of  $\text{C}_{1-6}$ -alk(en/yn)yl,  $\text{C}_{3-8}$ -cycloalk(en)yl and  $\text{C}_{3-8}$ -cycloalk(en)yl- $\text{C}_{1-6}$ -alk(en/yn)yl.

Yet another embodiment of the invention relates to compounds of formula I, wherein  
20  $\text{R}^2$  is selected from the group consisting of Ar, Ar- $\text{C}_{1-6}$ -alk(en/yn)yl, Ar- $\text{C}_{3-8}$ -cycloalk(en)yl and Ar- $\text{C}_{3-8}$ -cycloalk(en)yl- $\text{C}_{1-6}$ -alk(en/yn)yl.

Yet another embodiment of the invention relates to compounds of formula I, wherein  $\text{R}^2$  is selected from the group consisting of halogen, halo- $\text{C}_{1-6}$ -alk(en/yn)yl, halo- $\text{C}_{3-8}$ -  
25 cycloalk(en)yl, halo- $\text{C}_{3-8}$ -cycloalk(en)yl- $\text{C}_{1-6}$ -alk(en/yn)yl and cyano;  
with the proviso that when  $\text{R}^2$  is halogen or cyano then s is 0;.

In a preferred embodiment, the invention relates to compounds of formula I, wherein  $\text{R}^2$  is  $\text{NO}_2$  or a hydrogen atom;  
30 with the proviso that when  $\text{R}^2$  is  $\text{NO}_2$  then s is 0; and  
with the proviso that when  $\text{R}^2$  is a hydrogen atom and s is 1 then U is  $\text{NR}^{11}$ , O or S.

In one embodiment, the invention relates to compounds of formula I, wherein  $R^2$  is  $NO_2$  or a hydrogen atom or a halogen atom;  
with the proviso that when  $R^2$  is  $NO_2$  or a halogen atom then s is 0; and  
with the proviso that when  $R^2$  is a hydrogen atom and s is 1 then U is  $NR^{11}$ , O or S.

5

In another embodiment, the invention relates to compounds of formula I, wherein s is 0 and  $R^2$  is  $NO_2$  or a halogen atom.

10

In one embodiment, the invention relates to compounds of formula I, wherein  $R^2$  is a hydrogen atom.

In one embodiment, the invention relates to compounds of formula I, wherein s is 0 and  $R^2$  is selected from the group consisting of  $NO_2$ , halogen and cyano.

15 In another embodiment, the invention relates to compounds of formula I, wherein s is 0 and  $R^2$  is a hydrogen atom.

In yet another embodiment, the invention relates to compounds of formula I, wherein  $R^2$  is  $C_{1-6}$ -alk(en/yn)yl, typically  $C_{1-3}$ -alk(en/yn)yl.

20

In yet another embodiment, the invention relates to compounds of formula I, wherein  $R^2$  is  $C_{3-8}$ -cycloalk(en)yl, typically  $C_{3-6}$ -cycloalk(en)yl.

25

In yet another embodiment, the invention relates to compounds of formula I, wherein  $R^2$  is Ar.

In yet another embodiment, the invention relates to compounds of formula I, wherein  $R^2$  is Ar- $C_{1-6}$ -alk(en/yn)yl, typically Ar- $C_{1-3}$ -alk(en/yn)yl.

30 In yet another embodiment, the invention relates to compounds of formula I, wherein  $R^2$  is halo- $C_{1-6}$ -alk(en/yn)yl, typically halo- $C_{1-3}$ -alk(en/yn)yl.

In yet another embodiment, the invention relates to compounds of formula I, wherein  $s$  is 0 and  $R^2$  is a halogen atom.

5 In yet another embodiment, the invention relates to compounds of formula I, wherein  $s$  is 0 and  $R^2$  is cyano.

In another preferred embodiment, the invention relates to compounds of formula I, wherein  $s$  is 0 and  $R^2$  is  $NO_2$ .

10 In a preferred embodiment, the invention relates to compounds of formula I, wherein  $R^2$  is a hydrogen atom; with the proviso that when  $s$  is 1 then  $U$  is  $NR^{11}$ , O or S.

In one embodiment, the invention relates to compounds of formula I, wherein  $R^2$  is a hydrogen atom; with the proviso that when  $s$  is 1 then  $U$  is  $NR^{11}$ .

15

In another embodiment, the invention relates to compounds of formula I, wherein  $R^2$  is a hydrogen atom,  $s$  is 1,  $U$  is  $NR^{11}$  and  $R^{11}$  is a hydrogen atom.

20 In one embodiment, the invention relates to compounds of formula I, wherein the group  $-(U)_s-R^2$  is linked to position 6 of the indole or indoline.

In a preferred embodiment, the invention relates to compounds of formula I, wherein the group  $-(U)_s-R^2$  is linked to position 4 of the indole or indoline.

25 In a preferred embodiment, the invention relates to compounds of formula I, wherein  $X$  is CO.

In a preferred embodiment, the invention relates to compounds of formula I, wherein  $X$  is  $SO_2$ .

30

In a preferred embodiment, the invention relates to compounds of formula I, wherein  $q$  is 0.

In a preferred embodiment, the invention relates to compounds of formula **I**, wherein **q** is 1.

- 5 In one embodiment, the invention relates to compounds of formula **I**, wherein **q** is 1 and **Z** is a sulphur atom.

In a preferred embodiment, the invention relates to compounds of formula **I**, wherein **q** is 1 and **Z** is an oxygen atom.

10

In one embodiment, the invention relates to compounds of formula **I**, wherein **X** is SO<sub>2</sub> and **q** is 0.

15

In one embodiment, the invention relates to compounds of formula **I**, wherein **X** is CO and **q** is 0.

In one embodiment, the invention relates to compounds of formula **I**, wherein **X** is CO, **q** is 1 and **Z** is an oxygen atom.

- 20 In one embodiment, the invention relates to compounds of formula **I**, wherein **R**<sup>3</sup> is selected from the group consisting of C<sub>1-6</sub>-alk(en/yn)yl-heterocycloalk(en)yl, Ar-heterocycloalk(en)yl, Ar-C<sub>1-6</sub>-alk(en/yn)yl-heterocycloalk(en)yl, C<sub>1-6</sub>-alk(en/yn)yl-oxy-heterocycloalk(en)yl, C<sub>1-6</sub>-alk(en/yn)yl-oxy-carbonyl-C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl-oxy-carbonyl-C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl-oxy-carbonyl-C<sub>1-6</sub>-alk(en/yn)yl, hydroxy-C<sub>1-6</sub>-alk(en/yn)yl, hydroxy-C<sub>3-8</sub>-cycloalk(en)yl, hydroxy-heterocycloalk(en)yl, hydroxy-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, hydroxy-C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl, hydroxy-C<sub>1-6</sub>-alk(en/yn)yl-heterocycloalk(en)yl, halo-heterocycloalk(en)yl, halo-C<sub>1-6</sub>-alk(en/yn)yl-heterocycloalk(en)yl, cyano-C<sub>1-6</sub>-alk(en/yn)yl, cyano-C<sub>3-8</sub>-cycloalk(en)yl, cyano-heterocycloalk(en)yl, cyano-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, cyano-C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl, cyano-C<sub>1-6</sub>-alk(en/yn)yl-heterocycloalk(en)yl, acyl-C<sub>1-6</sub>-alk(en/yn)yl, acyl-C<sub>3-8</sub>-cycloalk(en)yl, acyl-heterocycloalk(en)yl, acyl-C<sub>3-8</sub>-
- 25
- 30



cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, acyl-C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl and acyl-C<sub>1-6</sub>-alk(en/yn)yl-heterocycloalk(en)yl.

In another embodiment, the invention relates to compounds of formula I, wherein **R**<sup>3</sup> is selected from the group consisting of C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl, heterocycloalk(en)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl, Ar, Ar-C<sub>1-6</sub>-alk(en/yn)yl, Ar-C<sub>3-8</sub>-cycloalk(en)yl, Ar-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, Ar-C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl, C<sub>1-6</sub>-alk(en/yn)yl-oxy-C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl-oxy-C<sub>1-6</sub>-alk(en/yn)yl, C<sub>1-6</sub>-alk(en/yn)yl-oxy-C<sub>3-8</sub>-cycloalk(en)yl, Ar-oxy-C<sub>1-6</sub>-alk(en/yn)yl, Ar-C<sub>1-6</sub>-alk(en/yn)yl-oxy-C<sub>1-6</sub>-alk(en/yn)yl, halo-C<sub>1-6</sub>-alk(en/yn)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, halo-C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl, halo-C<sub>1-6</sub>-alk(en/yn)yl-Ar, halo-C<sub>3-8</sub>-cycloalk(en)yl-Ar, halo-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl-Ar, halo-C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl-Ar and -NR<sup>12</sup>R<sup>12'</sup>; with the proviso that when **R**<sup>3</sup> is NR<sup>12</sup>R<sup>12'</sup> then q is 0.

In yet another embodiment, the invention relates to compounds of formula I, wherein **R**<sup>3</sup> is selected from the group consisting of C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl, heterocycloalk(en)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl, Ar, Ar-C<sub>1-6</sub>-alk(en/yn)yl, Ar-C<sub>3-8</sub>-cycloalk(en)yl, Ar-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, Ar-C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl, Ar-oxy-C<sub>1-6</sub>-alk(en/yn)yl, Ar-C<sub>1-6</sub>-alk(en/yn)yl-oxy-C<sub>1-6</sub>-alk(en/yn)yl, halo-C<sub>1-6</sub>-alk(en/yn)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, halo-C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl and -NR<sup>12</sup>R<sup>12'</sup>; with the proviso that when **R**<sup>3</sup> is NR<sup>12</sup>R<sup>12'</sup> then q is 0.

In yet another embodiment, the invention relates to compounds of formula I, wherein **R**<sup>3</sup> is selected from the group consisting of C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl, Ar, Ar-C<sub>1-6</sub>-alk(en/yn)yl, Ar-C<sub>3-8</sub>-cycloalk(en)yl, Ar-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, Ar-C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl, Ar-oxy-C<sub>1-6</sub>-alk(en/yn)yl, Ar-C<sub>1-6</sub>-alk(en/yn)yl-oxy-C<sub>1-6</sub>-alk(en/yn)yl and -NR<sup>12</sup>R<sup>12'</sup>; with the proviso that when **R**<sup>3</sup> is NR<sup>12</sup>R<sup>12'</sup> then q is 0.

In a preferred embodiment, the invention relates to compounds of formula I, wherein  $R^3$  is selected from the group consisting of  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl, heterocycloalk(en)yl, Ar, Ar- $C_{1-6}$ -alk(en/yn)yl, Ar-oxy- $C_{1-6}$ -alk(en/yn)yl, Ar- $C_{1-6}$ -alk(en/yn)yloxy- $C_{1-6}$ -alk(en/yn)yl, halo- $C_{1-6}$ -alk(en/yn)yl and  $-NR^{12}R^{12'}$ ;

5 with the proviso that when  $R^3$  is  $NR^{12}R^{12'}$  then q is 0.

In another preferred embodiment, the invention relates to compounds of formula I, wherein  $R^3$  is selected from the group consisting of  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl, Ar, Ar- $C_{1-6}$ -alk(en/yn)yl, Ar-oxy- $C_{1-6}$ -alk(en/yn)yl, Ar- $C_{1-6}$ -alk(en/yn)yloxy- $C_{1-6}$ -alk(en/yn)yl and  $-NR^{12}R^{12'}$ ;

10 with the proviso that when  $R^3$  is  $NR^{12}R^{12'}$  then q is 0.

In another preferred embodiment, the invention relates to compounds of formula I, wherein  $R^3$  is  $C_{1-6}$ -alk(en/yn)yl, typically  $C_{1-3}$ -alk(en/yn)yl.

15

In yet another preferred embodiment, the invention relates to compounds of formula I, wherein  $R^3$  is  $C_{3-8}$ -cycloalk(en)yl.

In yet another preferred embodiment, the invention relates to compounds of formula I, wherein  $R^3$  is  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl.

20

In yet another preferred embodiment, the invention relates to compounds of formula I, wherein  $R^3$  is heterocycloalk(en)yl.

25 In yet another preferred embodiment, the invention relates to compounds of formula I, wherein  $R^3$  is Ar.

In yet another preferred embodiment, the invention relates to compounds of formula I, wherein  $R^3$  is Ar- $C_{1-6}$ -alk(en/yn)yl.

30

In yet another preferred embodiment, the invention relates to compounds of formula I, wherein  $R^3$  is  $C_{1-6}$ -alk(en/yn)yl-oxy- $C_{1-6}$ -alk(en/yn)yl.

In yet another preferred embodiment, the invention relates to compounds of formula I, wherein  $R^3$  is Ar-oxy-C<sub>1-6</sub>-alk(en/yn)yl.

In yet another preferred embodiment, the invention relates to compounds of formula I,  
5 wherein  $R^3$  is Ar-C<sub>1-6</sub>-alk(en/yn)yloxy-C<sub>1-6</sub>-alk(en/yn)yl.

In yet another embodiment, the invention relates to compounds of formula I, wherein  $R^3$  is C<sub>1-6</sub>-alk(en/yn)yloxy-carbonyl-C<sub>1-6</sub>-alk(en/yn)yl.

10 In yet another embodiment, the invention relates to compounds of formula I, wherein  $R^3$  is halo-C<sub>1-6</sub>-alk(en/yn)yl, such as halo-C<sub>1-3</sub>-alk(en/yn)yl.

In yet another embodiment, the invention relates to compounds of formula I, wherein  $R^3$  is halo-C<sub>1-6</sub>-alk(en/yn)yl-Ar, such as halo-C<sub>1-3</sub>-alk(en/yn)yl-Ar.

15

In yet another preferred embodiment, the invention relates to compounds of formula I, wherein  $R^3$  is -NR<sup>12</sup>R<sup>12'</sup>, and q is 0.

In one embodiment, the invention relates to compounds of formula I, wherein X is  
20 CO, q is 1, Z is an oxygen atom and  $R^3$  is selected from the group consisting of C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl, Ar-C<sub>1-6</sub>-alk(en/yn)yl, Ar-C<sub>3-8</sub>-cycloalk(en)yl, Ar-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, Ar-C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl, Ar-oxy-C<sub>1-6</sub>-alk(en/yn)yl, Ar-C<sub>1-6</sub>-alk(en/yn)yloxy-C<sub>1-6</sub>-alk(en/yn)yl, halo-C<sub>1-6</sub>-alk(en/yn)yl,  
25 halo-C<sub>3-8</sub>-cycloalk(en)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl and halo-C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl.

In another embodiment, the invention relates to compounds of formula I, wherein X is CO, q is 1, Z is an oxygen atom and  $R^3$  is selected from the group consisting of C<sub>1-6</sub>-alk(en/yn)yl, Ar-C<sub>1-6</sub>-alk(en/yn)yloxy-C<sub>1-6</sub>-alk(en/yn)yl and halo-C<sub>1-6</sub>-alk(en/yn)yl.  
30

In one further embodiment, the invention relates to compounds of formula I, wherein X is CO, q is 0 and  $R^3$  is selected from the group consisting of C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-

cycloalk(en)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl, heterocycloalk(en)yl, Ar, Ar-C<sub>1-6</sub>-alk(en/yn)yl, Ar-C<sub>3-8</sub>-cycloalk(en)yl, Ar-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, Ar-C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl, Ar-oxy-C<sub>1-6</sub>-alk(en/yn)yl, Ar-C<sub>1-6</sub>-alk(en/yn)yl-oxy-C<sub>1-6</sub>-alk(en/yn)yl and -NR<sup>12</sup>R<sup>12'</sup>.

5

In yet another embodiment, the invention relates to compounds of formula I, wherein X is CO, q is 0 and R<sup>3</sup> is selected from the group consisting of C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl, heterocycloalk(en)yl, Ar, Ar-C<sub>1-6</sub>-alk(en/yn)yl, Ar-oxy-C<sub>1-6</sub>-alk(en/yn)yl and -NR<sup>12</sup>R<sup>12'</sup>.

10

In yet another embodiment, the invention relates to compounds of formula I, wherein X is SO<sub>2</sub>, q is 0 and R<sup>3</sup> is selected from the group consisting of C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl, Ar-C<sub>1-6</sub>-alk(en/yn)yl, Ar-C<sub>3-8</sub>-cycloalk(en)yl, Ar-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl and Ar-C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl.

15

In yet another embodiment, the invention relates to compounds of formula I, wherein X is SO<sub>2</sub>, q is 0 and R<sup>3</sup> is C<sub>1-6</sub>-alk(en/yn)yl or Ar-C<sub>1-6</sub>-alk(en/yn)yl.

20

In one embodiment, the invention relates to compounds of formula I, wherein R<sup>3</sup> is NR<sup>12</sup>R<sup>12'</sup> and q is 0 and wherein R<sup>12</sup> and R<sup>12'</sup> are independently selected from the group consisting of hydrogen, C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, Ar, Ar-C<sub>1-6</sub>-alk(en/yn)yl, Ar-C<sub>3-8</sub>-cycloalk(en)yl and Ar-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl.

25

In another embodiment, the invention relates to compounds of formula I, wherein R<sup>3</sup> is NR<sup>12</sup>R<sup>12'</sup> and q is 0 and wherein R<sup>12</sup> and R<sup>12'</sup> are independently selected from the group consisting of hydrogen, C<sub>1-6</sub>-alk(en/yn)yl, Ar and Ar-C<sub>1-6</sub>-alk(en/yn)yl or wherein R<sup>12</sup> and R<sup>12'</sup> together with the nitrogen atom to which they are attached form a 4-8 membered saturated or unsaturated ring which optionally contains 1, 2 or 3 further heteroatoms.

30

In a preferred embodiment, the invention relates to compounds of formula I, wherein  $R^3$  is  $NR^{12}R^{12'}$  and  $q$  is 0 and wherein  $R^{12}$  and  $R^{12'}$  are independently selected from the group consisting of hydrogen,  $C_{1-6}$ -alk(en/yn)yl, Ar and Ar- $C_{1-6}$ -alk(en/yn)yl.

- 5 In another preferred embodiment, the invention relates to compounds of formula I, wherein  $R^3$  is  $NR^{12}R^{12'}$  and  $q$  is 0 and wherein  $R^{12}$  and  $R^{12'}$  together with the nitrogen atom to which they are attached form a 4-8 membered saturated or unsaturated ring which optionally contains 1, 2 or 3 further heteroatoms.
- 10 In one embodiment, the invention relates to compounds of formula I, wherein  $R^3$  is  $NR^{12}R^{12'}$  and  $q$  is 0 and wherein at least one of  $R^{12}$  and  $R^{12'}$  is a hydrogen atom.

- In another embodiment, the invention relates to compounds of formula I, wherein  $R^3$  is  $NR^{12}R^{12'}$  and  $q$  is 0 and at least one of  $R^{12}$  and  $R^{12'}$  is  $C_{1-6}$ -alk(en/yn)yl, typically
- 15  $C_{1-3}$ -alk(en/yn)yl.

In yet another embodiment, the invention relates to compounds of formula I, wherein  $R^3$  is  $NR^{12}R^{12'}$  and  $q$  is 0 and one of  $R^{12}$  and  $R^{12'}$  is Ar.

- 20 In yet another embodiment, the invention relates to compounds of formula I, wherein  $R^3$  is  $NR^{12}R^{12'}$  and  $q$  is 0 and one of  $R^{12}$  and  $R^{12'}$  is Ar- $C_{1-6}$ -alk(en/yn)yl, typically Ar- $C_{1-3}$ -alk(en/yn)yl.

- In a preferred embodiment, the invention relates to compounds of formula I, wherein
- 25 Y is of formula II, III, V, XXX or XXXI.

In one embodiment, the invention relates to compounds of formula I, wherein Y is of formula III or IV.

- 30 In a preferred embodiment, the invention relates to compounds of formula I, wherein Y is of formula II or V.

In a preferred embodiment, the invention relates to compounds of formula I, wherein Y is of formula V or XXXI.

In a preferred embodiment, the invention relates to compounds of formula I, wherein  
5 Y is of formula II or III and W is a sulphur atom.

In a preferred embodiment, the invention relates to compounds of formula I, wherein Y is of formula II or III and W is an oxygen atom.

10 In a preferred embodiment, the invention relates to compounds of formula I, wherein Y is of formula V.

In a preferred embodiment, the invention relates to compounds of formula I, wherein Y is of formula XXX and T is NH.

15 In a preferred embodiment, the invention relates to compounds of formula I, wherein Y is of formula XXX and T is a nitrogen atom or an oxygen atom.

In a preferred embodiment, the invention relates to compounds of formula I, wherein  
20 Y is of formula XXXI and L is a nitrogen atom.

In a preferred embodiment, the invention relates to compounds of formula I, wherein Y is of formula XXXI and L is C or CH.

25 In one embodiment, the invention relates to compounds of formula I, wherein each  $R^5$  is independently selected from the group consisting of Ar-C<sub>1-6</sub>-alk(en/yn)yl, acyl, -CO-NR<sup>6</sup>R<sup>6'</sup>, cyano, cyano-C<sub>1-6</sub>-alk(en/yn)yl, cyano-C<sub>3-8</sub>-cycloalk(en)yl and cyano-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl.

30 In another embodiment, the invention relates to compounds of formula I, wherein each  $R^5$  is independently selected from the group consisting of C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, Ar, C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, halogen, halo-C<sub>1-6</sub>-

alk(en/yn)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, -NR<sup>7</sup>R<sup>7'</sup>, -S-R<sup>8</sup> and -SO<sub>2</sub>R<sup>8</sup>; or

two adjacent R<sup>5</sup> together with the aromatic group to which they are attached form a 4-8 membered ring which optionally contains one or two heteroatoms.

5

In yet another embodiment, the invention relates to compounds of formula I, wherein each R<sup>5</sup> is independently selected from the group consisting of C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, Ar, C<sub>1-6</sub>-alk(en/yn)oxy, C<sub>3-8</sub>-cycloalk(en)oxy, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)oxy, -NR<sup>7</sup>R<sup>7'</sup>, -S-R<sup>8</sup> and -

10 SO<sub>2</sub>R<sup>8</sup>; or

two adjacent R<sup>5</sup> together with the aromatic group to which they are attached form a 4-8 membered ring which optionally contains one or two heteroatoms.

In yet another embodiment, the invention relates to compounds of formula I, wherein

15 each R<sup>5</sup> is independently selected from the group consisting of halogen, halo-C<sub>1-6</sub>-alk(en/yn)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl and halo-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl.

In yet another embodiment, the invention relates to compounds of formula I, wherein each R<sup>5</sup> is independently selected from the group consisting of C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, Ar, Ar-thio, Ar-oxy, halogen, halo-C<sub>1-6</sub>-alk(en/yn)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, or

20

two adjacent R<sup>5</sup> together with the aromatic group to which they are attached form a 4-8 membered ring which optionally contains one or two heteroatoms.

25

In a preferred embodiment, the invention relates to compounds of formula I, wherein each R<sup>5</sup> is independently selected from the group consisting of C<sub>1-6</sub>-alk(en/yn)yl, Ar, Ar-thio, Ar-oxy, halogen, halo-C<sub>1-6</sub>-alk(en/yn)yl, or

two adjacent R<sup>5</sup> together with the aromatic group to which they are attached form a 4-

30 8 membered ring which optionally contains one or two heteroatoms.

In another preferred embodiment, the invention relates to compounds of formula I, wherein each  $R^5$  is independently selected from the group consisting of halogen and halo-C<sub>1-6</sub>-alk(en/yn)yl.

- 5 In an embodiment, the invention relates to compounds of formula I, wherein at least one substituent  $R^5$  is a halogen atom.

In another embodiment, the invention relates to compounds of formula I, wherein at least one substituent  $R^5$  is halo-C<sub>1-6</sub>-alk(en/yn)yl, typically halo-C<sub>1-3</sub>-alk(en/yn)yl.

10

In yet another embodiment, the invention relates to compounds of formula I, wherein at least one substituent  $R^5$  is C<sub>1-6</sub>-alk(en/yn)yl.

15

In yet another embodiment, the invention relates to compounds of formula I, wherein at least one substituent  $R^5$  is Ar.

In yet another embodiment, the invention relates to compounds of formula I, wherein at least one substituent  $R^5$  is Ar-thio.

20

In yet another embodiment, the invention relates to compounds of formula I, wherein at least one substituent  $R^5$  is Ar-oxy.

In yet another embodiment, the invention relates to compounds of formula I, wherein at least one substituent  $R^5$  is C<sub>1-6</sub>-alk(en/yn)yloxy.

25

In yet another embodiment, the invention relates to compounds of formula I, wherein at least one substituent  $R^5$  is -NR<sup>7</sup>R<sup>7'</sup>.

30

In yet another embodiment, the invention relates to compounds of formula I, wherein at least one substituent  $R^5$  is -S-R<sup>8</sup>.

In yet another embodiment, the invention relates to compounds of formula I, wherein at least one substituent  $R^5$  is -SO<sub>2</sub>R<sup>8</sup>.



In yet another embodiment, the invention relates to compounds of formula I, wherein two adjacent  $R^5$  together with the aromatic group form a 4-8 membered ring, which optionally contains one or two heteroatoms.

- 5 In a preferred embodiment, the invention relates to compounds of formula I, wherein two adjacent  $R^5$  together form

$-(CH_2)_n-CH_2-$ ,  $-CH=CH-(CH_2)_{m'}-$ ,  $-CH_2-CH=CH-(CH_2)_{p'}-$ ,  $-CH=CH-CH=CH-$ ,

$-(CH_2)_n-O-$ ,  $-O-(CH_2)_{m'}-O-$ ,  $-CH_2-O-(CH_2)_{p'}-O-$ ,  $-CH_2-O-CH_2-O-CH_2-$ ,

$-(CH_2)_n-S-$ ,  $-S-(CH_2)_{m'}-S-$ ,  $-CH_2-S-(CH_2)_{p'}-S-$ ,  $-CH_2-S-CH_2-S-CH_2-$ ,

- 10  $-(CH_2)_n-NH-$ ,  $-NH-(CH_2)_{m'}-NH-$ ,  $-CH_2-NH-(CH_2)_{p'}-NH-$ ,  $-CH=CH-NH-$ ,  
 $-O-(CH_2)_{m'}-NH-$ ,  $-CH_2-O-(CH_2)_{p'}-NH-$  or  $-O-(CH_2)_{p'}-NH-CH_2-$ ,  $-S-(CH_2)_{m'}-NH-$ ,  
 $-N=CH-NH-$ ,  $-N=CH-O-$  or  $-N=CH-S-$ , wherein  $m'$  is 1, 2 or 3,  $n'$  is 2, 3 or 4 and  $p'$  is 1 or 2.

- 15 In yet another embodiment, the invention relates to compounds of formula I, wherein two adjacent  $R^5$  together form  $-CH_2-O-CH_2-$ .

In yet another embodiment, the invention relates to compounds of formula I, wherein two adjacent  $R^5$  together form  $-CH=CH-CH=CH-$ .

20

In yet another embodiment, the invention relates to compounds of formula I, wherein two adjacent  $R^5$  together form  $-O-CH_2-O-$ .

- 25 In yet another embodiment, the invention relates to compounds of formula I, wherein two adjacent  $R^5$  together form  $-O-CH_2-O-CH_2-$ .

- In yet another embodiment, the invention relates to compounds of formula I, wherein at least one substituent  $R^5$  is  $-NR^7R^{7'}$ ; and wherein  $R^7$  and  $R^{7'}$  are independently selected from the group consisting of hydrogen,  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl  
 30 and  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl.

In yet another embodiment, the invention relates to compounds of formula **I**, wherein at least one substituent  $R^5$  is  $-NR^7R^{7'}$ ; and wherein  $R^7$  and  $R^{7'}$  are independently selected from the group consisting of hydrogen and  $C_{1-6}$ -alk(en/yn)yl.

- 5 In yet another embodiment, the invention relates to compounds of formula **I**, wherein at least one substituent  $R^5$  is  $-NR^7R^{7'}$ ; and wherein both  $R^7$  and  $R^{7'}$  are  $C_{1-6}$ -alk(en/yn)yl, typically  $C_{1-3}$ -alk(en/yn)yl.
- 10 In yet another embodiment, the invention relates to compounds of formula **I**, wherein at least one substituent  $R^5$  is  $-S-R^8$  or  $-SO_2R^8$ ; and wherein  $R^8$  is selected from the group consisting of hydrogen,  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl and Ar .
- 15 In yet another embodiment, the invention relates to compounds of formula **I**, wherein at least one substituent  $R^5$  is  $-S-R^8$  or  $-SO_2R^8$ ; and wherein  $R^8$  is selected from the group consisting of  $C_{1-6}$ -alk(en/yn)yl and Ar .
- 20 One embodiment of the invention relates to compounds of formula **I**, wherein  $s$  is 0 and  $q$  is 0.

Another embodiment of the invention relates to compounds of formula **I**, wherein  $R^2$  is a hydrogen atom and  $X$  is CO.

25

Yet another embodiment of the invention relates to compounds of formula **I**, wherein  $s$  is 0 and  $X$  is CO.

- Yet another embodiment of the invention relates to compounds of formula **I**, wherein  $R^2$  is a hydrogen atom and  $q$  is 0.
- 30

Yet another embodiment of the invention relates to compounds of formula **I**, wherein  $q$  is 0 and  $X$  is CO.

One embodiment of the invention relates to compounds of formula I, wherein the total number of Ar-groups in the substituents  $R^2$ ,  $R^3$  and  $R^5$  equals to 0, 1, 2, or 3, typically 0 or 1.

- 5 Another embodiment of the invention relates to compounds of formula I wherein neither  $R^2$ ,  $R^3$  or  $R^5$  comprises an Ar-group.

Yet another embodiment of the invention relates to compounds of formula I, wherein the total number of Ar-groups in the substituents  $R^2$ ,  $R^3$  and  $R^5$  equals to 1.

10

Yet another embodiment of the invention relates to compounds of formula I, wherein the total number of Ar-groups in the substituents  $R^2$ ,  $R^3$  and  $R^5$  equals to 2.

- One embodiment of the invention relates to compounds of formula I, wherein  $R^3$  is not  $CH_3$  when X is  $SO_2$  and q is 0.

15

Another embodiment of the invention relates to compounds of formula I, wherein  $X-(Z)_q-R^3$  is not  $SO_2-CH_3$  when Y is of formula V.

- 20 Yet another embodiment of the invention relates to compounds of formula I, wherein  $R^3$  is  $NR^{12}R^{12'}$  and both  $R^{12}$  and  $R^{12'}$  is different from Ar.

- Yet another embodiment of the invention relates to compounds of formula I, wherein  $R^3$  is  $NR^{12}R^{12'}$  and one of  $R^{12}$  and  $R^{12'}$  is Ar, with the proviso that Ar is different from quinoline or phenyl.

25

Another embodiment of the invention relates to compounds of formula I, wherein Y is not of formula V when X is CO and q is 0 and  $R^3$  is  $NR^{12}R^{12'}$  and one of  $R^{12}$  and  $R^{12'}$  is Ar, typically quinoline or phenyl.

30

In another embodiment, the compound of formula I is not:

N-[1-(phenylmethyl)-1H-indol-5-yl]-Methanesulfonamide;

N-[1-[(4-fluorophenyl)methyl]-1H-indol-5-yl]-Methanesulfonamide;

N-[2,3-dihydro-1-(phenylmethyl)-1H-indol-5-yl]-Methanesulfonamide;


N-[1-(phenylmethyl)-1H-indol-5-yl]-N'-4-quinoliny-Urea;

N-[1-(phenylmethyl)-1H-indol-5-yl]-N'-4-quinoliny-Urea; or

1-(1-benzyl-5-indoliny)-3-phenyl-Urea.

5

One aspect of the invention, relates to compounds of general formula **VII** and salts thereof:



10 wherein the dotted line, **f**, **q**, **s**, **U**, **X**, **Z**, **R<sup>1</sup>**, **R<sup>1'</sup>**, **R<sup>2</sup>**, **R<sup>3</sup>** and **R<sup>5</sup>** are as defined under formula **I**. Any of the embodiments related to formula **I** are also embodiments of formula **VII**.

15 In one embodiment, the invention relates to compounds of the general formula **VII**, wherein **f** is 0.

In another embodiment, the invention relates to compounds of the general formula **VII** being substituted by one substituent **R<sup>5</sup>**, such as in the orto-, meta- or para-position.


20

In a preferred embodiment, the invention relates to compounds of the general formula **VII**, which are substituted by one substituent **R<sup>5</sup>** in the para-position.

In one embodiment, the invention relates to compounds of the general formula **VII** being substituted by two independently selected  $R^5$  substituents, such as in the ortho- and para-position, in the meta- and para-position and in the orto- and meta-position.

- 5 In another embodiment, the invention relates to compounds of the general formula **VII** being substituted by three independently selected  $R^5$  substituents.

Another aspect of the invention relates to compounds of the general formula **VIII** or  
10 salts thereof:



wherein the dotted line,  $g$ ,  $h$ ,  $q$ ,  $s$ ,  $U$ ,  $X$ ,  $Z$ ,  $R^1$ ,  $R^{1'}$ ,  $R^2$ ,  $R^3$  and  $R^5$  are as defined under  
15 formula **I**. Any of the embodiments related to formula **I** are also embodiments of  
formula **VIII**.

In one embodiment, the invention relates to compounds of the general formula **VIII**,  
wherein the nitrogen atom is attached to position 1 of the naphthyl group via the  
methylene group.

20

In another embodiment, the invention relates to compounds of the general formula  
**VIII**, wherein the nitrogen atom is attached to position 2 of the naphthyl group via the  
methylene group.

In yet another embodiment, the invention relates to compounds of the general formula **VIII**, wherein **g** is 0, 1, 2 or 3, typically 0, 1 or 2.

In yet another embodiment, the invention relates to compounds of the general formula **VIII**, wherein **h** is 0, 1 or 2, typically 0 or 1.

In yet another embodiment, the invention relates to compounds of the general formula **VIII**, wherein **g + h** equals to 0, 1, 2 or 3.


In yet another embodiment, the invention relates to compounds of the general formula **VIII**, wherein both **g** and **h** are 0.

In yet another embodiment, the invention relates to compounds of the general formula **VIII** being substituted by one substituent **R<sup>5</sup>**.

In yet another embodiment, the invention relates to compounds of the general formula **VIII** being substituted by two independently selected **R<sup>5</sup>** substituents.

In yet another embodiment, the invention relates to compounds of the general formula **VIII** being substituted by three independently selected **R<sup>5</sup>** substituents.

Yet another aspect of the invention relates to compounds of the general formula **IX** or salts thereof:



wherein the dotted line, **a**, **q**, **s**, **U**, **X**, **Z**, **R<sup>1</sup>**, **R<sup>1'</sup>**, **R<sup>2</sup>**, **R<sup>3</sup>** and **R<sup>5</sup>** are as defined under  
 formula I. Any of the embodiments related to formula I are also embodiments of  
 5 formula IX.

In an embodiment, the invention relates to compounds of the general formula IX,  
 wherein the nitrogen atom is attached to position 2 of the heteroaromatic group via the  
 methylene group.

In another embodiment, the invention relates to compounds of the general formula IX,  
 wherein the nitrogen atom is attached to position 3 of the heteroaromatic group via the  
 methylene group.

In yet another embodiment, the invention relates to compounds of the general formula  
 IX, wherein **W** is an oxygen atom.

In a preferred embodiment, the invention relates to compounds of the general formula  
 IX, wherein **W** is a sulphur atom.

In another embodiment, the invention relates to compounds of the general formula IX,  
 wherein **a** is 0, 1 or 2.


In yet another embodiment, the invention relates to compounds of the general formula IX, wherein **a** is 0.

In yet another embodiment, the invention relates to compounds of the general formula IX being substituted by one substituent **R**<sup>5</sup>, such as in position 5.

In yet another embodiment, the invention relates to compounds of the general formula IX being substituted by two independently selected **R**<sup>5</sup> substituents.

10 In an embodiment, the invention relates to compounds of the general formula IX, wherein the nitrogen atom is attached to position 2 via the methylene group and wherein a substituent **R**<sup>5</sup> is attached to position 5 of the heteroaromatic group.

15 Yet another aspect of the invention relates to compounds of the general formula X or salts thereof:



wherein the dotted line, **b**, **c**, **q**, **s**, **U**, **X**, **Z**, **R**<sup>1</sup>, **R**<sup>1'</sup>, **R**<sup>2</sup>, **R**<sup>3</sup> and **R**<sup>5</sup> are as defined under formula I. Any of the embodiments related to formula I are also embodiments of formula X.



In an embodiment, the invention relates to compounds of the general formula **X**, wherein the nitrogen atom is attached to position 2 of the heteroaromatic group via the methylene group.

- 5 In another embodiment, the invention relates to compounds of the general formula **X**, wherein the nitrogen atom is attached to position 3 of the heteroaromatic group via the methylene group.

- In yet another embodiment, the invention relates to compounds of the general formula  
10 **X**, wherein **W** is an oxygen atom.

In yet another embodiment, the invention relates to compounds of the general formula **X**, wherein **W** is a sulphur atom.

- 15 In yet another embodiment, the invention relates to compounds of the general formula **X**, wherein **b** is 0, 1, 2 or 3, typically 0, 1 or 2.

- In yet another embodiment, the invention relates to compounds of the general formula **X**, wherein **c** is 0 or 1, typically 0.  
20

In yet another embodiment, the invention relates to compounds of the general formula **X**, wherein **b + c** equals to 0, 1, 2, 3 or 4.

- In yet another embodiment, the invention relates to compounds of the general formula  
25 **X**, wherein both **b** and **c** are 0.


- In yet another embodiment, the invention relates to compounds of the general formula **X**, wherein **b + c** equals to 1. In one aspect thereof **b** is 1 and **c** is 0. In another aspect thereof **b** is 0 and **c** is 1.  
30

In yet another embodiment, the invention relates to compounds of the general formula **X** being substituted by one substituent **R**<sup>5</sup>.

In yet another embodiment, the invention relates to compounds of the general formula X being substituted by two independently selected  $R^5$  substituents.

In yet another embodiment, the invention relates to compounds of the general formula  
5 X being substituted by three independently selected  $R^5$  substituents.

Yet another aspect of the invention relates to compounds of the general formula XI or salts thereof:



wherein the dotted line, d, e, q, s, U, X, Z,  $R^1$ ,  $R^{1'}$ ,  $R^2$ ,  $R^3$  and  $R^5$  are as defined under formula I. Any of the embodiments related to formula I are also embodiments of formula XI.

In an embodiment, the invention relates to compounds of the general formula XI, wherein the nitrogen atom is attached to position 4 of the heteroaromatic group via the methylene group.

20 In another embodiment, the invention relates to compounds of the general formula XI, wherein the nitrogen atom is attached to position 5 of the heteroaromatic group via the methylene group.

In an embodiment, the invention relates to compounds of the general formula **XI**, wherein the nitrogen atom is attached to position 6 of the heteroaromatic group via the methylene group.

- 5 In another embodiment, the invention relates to compounds of the general formula **XI**, wherein the nitrogen atom is attached to position 7 of the heteroaromatic group via the methylene group.

- In yet another embodiment, the invention relates to compounds of the general formula  
10 **XI**, wherein **W** is an oxygen atom.

In yet another embodiment, the invention relates to compounds of the general formula **XI**, wherein **W** is a sulphur atom.

- 15 In yet another embodiment, the invention relates to compounds of the general formula **XI**, wherein **d** is 0, 1 or 2, typically 0 or 1.

In yet another embodiment, the invention relates to compounds of the general formula **XI**, wherein **e** is 0, 1 or 2.

20

In yet another embodiment, the invention relates to compounds of the general formula **XI**, wherein **d + e** is 0, 1, 2, 3 or 4.

- In yet another embodiment, the invention relates to compounds of the general formula  
25 **XI**, wherein both **d** and **e** are 0.


In yet another embodiment, the invention relates to compounds of the general formula **XI** being substituted by one substituent **R<sup>5</sup>**.

- 30 In yet another embodiment, the invention relates to compounds of the general formula **XI** being substituted by two independently selected **R<sup>5</sup>** substituents.

In yet another embodiment, the invention relates to compounds of the general formula **XI** being substituted by three independently selected  $R^5$  substituents.

Yet another aspect of the invention relates to compounds of the general formula

5 **XXXII** or salts thereof:



wherein the dotted line, **j**, **q**, **s**, **T**, **U**, **X**, **Z**, **R<sup>1</sup>**, **R<sup>1'</sup>**, **R<sup>2</sup>**, **R<sup>3</sup>** and **R<sup>5</sup>** are as defined under formula I. Any of the embodiments related to formula I are also embodiments of formula XXXII.

In an embodiment, the invention relates to compounds of the general formula **XXXII**, wherein the nitrogen atom is attached to the position indicated with 1 of the heteroaromatic group via the methylene group.

In another embodiment, the invention relates to compounds of the general formula **XXXII**, wherein the nitrogen atom is attached to the position indicated with 2 of the heteroaromatic group via the methylene group.

20 In another embodiment, the invention relates to compounds of the general formula  
**XXXII**, wherein the nitrogen atom is attached to the position indicated with 3 of the  
heteroaromatic group via the methylene group.

In yet another embodiment, the invention relates to compounds of the general formula **XXXII**, wherein **T** is an oxygen atom.

In yet another embodiment, the invention relates to compounds of the general formula **XXXII**, wherein **T** is a nitrogen atom.

In yet another embodiment, the invention relates to compounds of the general formula **XXXII**, wherein **T** represents NH.

In another embodiment, the invention relates to compounds of the general formula **XXXII**, wherein **j** is 0, 1, 2 or 3.

In yet another embodiment, the invention relates to compounds of the general formula **XXXII**, wherein **j** is 0.


In yet another embodiment, the invention relates to compounds of the general formula **XXXII** being substituted by at least one substituent **R<sup>5</sup>**. In one aspect thereof, the compound of the general formula **XXXII** is substituted in the position indicated with 1. In another aspect thereof, the compound of the general formula **XXXII** is substituted in the position indicated with 2. In yet another aspect thereof, the compound of the general formula **XXXII** is substituted in the position indicated with 3. In yet another aspect thereof, **T** represents a nitrogen atom at which the compound of the general formula **XXXII** is substituted.

In yet another embodiment, the invention relates to compounds of the general formula **XXXII** being substituted by two independently selected **R<sup>5</sup>** substituents.

In yet another embodiment, the invention relates to compounds of the general formula **XXXII** being substituted by two or three independently selected **R<sup>5</sup>** substituents.

In yet another embodiment, the invention relates to compounds of the general formula **XXXII** being substituted by three independently selected **R<sup>5</sup>** substituents.

One aspect of the invention, relates to compounds of general formula **XXXIII** and salts thereof:



- 5 wherein the dotted line, **k**, **q**, **s**, **L**, **U**, **X**, **Z**, **R<sup>1</sup>**, **R<sup>1'</sup>**, **R<sup>2</sup>**, **R<sup>3</sup>** and **R<sup>5</sup>** are as defined under formula I. Any of the embodiments related to formula I are also embodiments of formula **XXXIII**.

- 10 In one embodiment, the invention relates to compounds of the general formula **XXXIII**, wherein **k** is 0.

In another embodiment, the invention relates to compounds of the general formula **XXXIII** being substituted by one substituent **R<sup>5</sup>**, such as in the position being ortho, meta or para to the nitrogen atom.

15

In a preferred embodiment, the invention relates to compounds of the general formula **XXXIII**, which are substituted by one substituent **R<sup>5</sup>** in the position being para to the nitrogen atom.

- 20 In one embodiment, the invention relates to compounds of the general formula **XXXIII** being substituted by two independently selected **R<sup>5</sup>** substituents, such as in the in the positions being ortho and para to the nitrogen atom, or in the positions being meta and para to the nitrogen atom, or in the positions being ortho and meta to the nitrogen atom.

25

In another embodiment, the invention relates to compounds of the general formula **XXXIII** being substituted by three independently selected  $R^5$  substituents.

5

In one embodiment of the invention, the compounds of the following list and salts thereof are preferred:

*N*-[4-Chloro-1-(4-trifluoromethylbenzyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide,

10 *N*-[4-Chloro-1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide,

[1-(4-Fluorobenzyl)-2,3-dihydro-1H-indol-5-yl]-carbamic acid propyl ester,

*N*-[1-(4-Fluorobenzyl)-2,3-dihydro-1H-indol-5-yl]-*C*-phenyl-methanesulfonamide,

4-Fluoro-*N*-[1-(4-fluorobenzyl)-2,3-dihydro-1H-indol-5-yl]-benzamide,

15 *N*-[1-(4-Fluorobenzyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide,

*N*-[1-(4-Fluorobenzyl)-2,3-dihydro-1H-indol-5-yl]-2-thiophen-2-ylacetamide,

*N*-[1-(4-Fluorobenzyl)-2,3-dihydro-1H-indol-5-yl]-2-(4-fluorophenyl)-acetamide,

3-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-1,1-diisopropylurea,

Morpholine-4-carboxylic acid [1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-

20 indol-5-yl]-amide,

Pyrrolidine-1-carboxylic acid [1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-amide,

[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-carbamic acid 2-benzyloxyethyl ester,

25 3-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-1-methyl-1-propylurea,

[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-carbamic acid tert-butyl ester,

*N*-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-*C*-phenyl-

30 methanesulfonamide,

Butane-1-sulfonic acid [1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-amide,

*N*-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-4-fluorobenzamide,

- N*-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-2,2-dimethylpropionamide,  
*N*-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-2-phenoxyacetamide,  
5 *N*-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide,  
*N*-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-butyramide,  
Cyclopentanecarboxylic acid [1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-amide,  
10 *N*-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-2-thiophen-2-ylacetamide,  
*N*-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-isonicotinamide,  
*N*-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-4-dimethylaminobenzamide,  
15 *N*-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-2-(4-fluorophenyl)-acetamide,  
*N*-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-6-trifluoromethylnicotinamide,  
*1-tert-Butyl-3*-[1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-urea,  
20 *1*-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-3-ethylurea,  
*1-Benzyl-3*-[1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-urea,  
*1*-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-3-phenethylurea,  
*1*-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-3-thiophen-2-ylurea,  
*1*-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-3-thiophen-3-ylurea,  
25 *2,2-Dimethyl-N*-[6-nitro-1-(4-trifluoromethylbenzyl)-2,3-dihydro-1H-indol-5-yl]-propionamide,  
*N*-[1-(5-Chlorothiophen-2-ylmethyl)-6-nitro-2,3-dihydro-1H-indol-5-yl]-2,2-dimethylpropionamide,  
*2*-(4-Fluorophenyl)-*N*-[6-nitro-1-(4-trifluoromethylbenzyl)-2,3-dihydro-1H-indol-5-yl]-acetamide,  
30 *N*-[6-Amino-1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide,



- N*-[6-Amino-1-(4-trifluoromethylbenzyl)-2,3-dihydro-1*H*-indol-5-yl]-2,2-dimethylpropionamide,  
*N*-[6-Amino-1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-2,2-dimethylpropionamide,  
5 *N*-[6-Amino-1-(4-trifluoromethylbenzyl)-2,3-dihydro-1*H*-indol-5-yl]-2-(4-fluorophenyl)-acetamide, or  
*N*-[1-(5-Chlorothiophen-2-ylmethyl)-1*H*-indol-5-yl]-3,3-dimethylbutyramide.

10 In another embodiment of the invention, the compounds of the following list and salts thereof are preferred:

- N*-[4-Chloro-1-(4-trifluoromethylbenzyl)-2,3-dihydro-1*H*-indol-5-yl]-3,3-dimethylbutyramide;  
*N*-[4-Chloro-1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-3,3-dimethylbutyramide;  
15 [1-(4-Fluorobenzyl)-2,3-dihydro-1*H*-indol-5-yl]-carbamic acid propyl ester;  
*N*-[1-(4-Fluorobenzyl)-2,3-dihydro-1*H*-indol-5-yl]-*C*-phenyl-methanesulfonamide;  
4-Fluoro-*N*-[1-(4-fluorobenzyl)-2,3-dihydro-1*H*-indol-5-yl]-benzamide;  
*N*-[1-(4-Fluorobenzyl)-2,3-dihydro-1*H*-indol-5-yl]-3,3-dimethylbutyramide;  
*N*-[1-(4-Fluorobenzyl)-2,3-dihydro-1*H*-indol-5-yl]-2-thiophen-2-ylacetamide;  
20 *N*-[1-(4-Fluorobenzyl)-2,3-dihydro-1*H*-indol-5-yl]-2-(4-fluorophenyl)-acetamide;  
3-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-1,1-diisopropylurea;  
Morpholine-4-carboxylic acid [1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-amide;  
Pyrrolidine-1-carboxylic acid [1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-amide;  
25 [1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-carbamic acid 2-benzyloxyethyl ester;  
3-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-1-methyl-1-propylurea;  
30 [1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-carbamic acid tert-butyl ester;  
*N*-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-*C*-phenyl-methanesulfonamide;

- Butane-1-sulfonic acid [1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-amide;*
- N-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-4-fluorobenzamide;*
- N-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-2,2-*
- 5 *dimethylpropionamide;*
- N-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-2-phenoxyacetamide;*
- N-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide;*
- 10 *N-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-butyramide;*
- Cyclopentanecarboxylic acid [1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-amide;*
- N-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-2-thiophen-2-ylacetamide;*
- 15 *N-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-isonicotinamide;*
- N-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-4-dimethylaminobenzamide;*
- N-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-2-(4-fluorophenyl)-acetamide;*
- 20 *N-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-6-trifluoromethylnicotinamide;*
- 1-tert-Butyl-3-[1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-urea;*
- 1-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-3-ethylurea;*
- 1-Benzyl-3-[1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-urea;*
- 25 *1-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-3-phenethylurea;*
- 1-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-3-thiophen-2-ylurea;*
- 1-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-3-thiophen-3-ylurea;*
- [1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-carbamic acid propyl ester;*
- 30 *2,2-Dimethyl-N-[6-nitro-1-(4-trifluoromethylbenzyl)-2,3-dihydro-1H-indol-5-yl]-propionamide;*
- N-[1-(5-Chlorothiophen-2-ylmethyl)-6-nitro-2,3-dihydro-1H-indol-5-yl]-2,2-dimethylpropionamide;*

- 2-(4-Fluorophenyl)-N-[6-nitro-1-(4-trifluoromethylbenzyl)-2,3-dihydro-1H-indol-5-yl]-acetamide;
- N-[1-(5-Chlorothiophen-2-ylmethyl)-6-nitro-2,3-dihydro-1H-indol-5-yl]-2-(4-fluorophenyl)-acetamide;
- 5 N-[1-(5-Chlorothiophen-2-ylmethyl)-6-nitro-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide;
- N-[6-Amino-1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide;
- N-[6-Amino-1-(4-trifluoromethylbenzyl)-2,3-dihydro-1H-indol-5-yl]-2,2-
- 10 dimethylpropionamide;
- N-[6-Amino-1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-2,2-dimethylpropionamide;
- N-[6-Amino-1-(4-trifluoromethylbenzyl)-2,3-dihydro-1H-indol-5-yl]-2-(4-fluorophenyl)-acetamide;
- 15 N-[6-Amino-1-(4-trifluoromethylbenzyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide;
- N-[6-Amino-1-(4-fluorobenzyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide;
- N-[6-Amino-1-(3-fluoro-4-trifluoromethylbenzyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide;
- 20 N-[1-(5-Chlorothiophen-2-ylmethyl)-1H-indol-5-yl]-3,3-dimethylbutyramide;
- N-[6-Bromo-1-(4-trifluoromethylbenzyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide;
- N-[6-Bromo-1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide;
- 25 N-[1-(4-Chlorobenzyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide;
- 3,3-Dimethyl-N-[1-(4-trifluoromethylbenzyl)-2,3-dihydro-1H-indol-5-yl]-butyramide;
- N-[1-(4-Isopropylbenzyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide;
- N-[1-(3-Fluoro-4-trifluoromethylbenzyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide;
- 30 N-[1-(6-Chlorobenzo[1,3]dioxol-5-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide;
- N-[1-(3,5-Dimethyl-1-phenyl-1H-pyrazol-4-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide;

- N*-[1-(2-Chloro-5-trifluoromethylbenzyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide;
- N*-{1-[5-(4-Chlorophenoxy)-1,3-dimethyl-1H-pyrazol-4-ylmethyl]-2,3-dihydro-1H-indol-5-yl}-3,3-dimethylbutyramide;
- 5    3,3-Dimethyl-*N*-[1-(6-*p*-tolylloxy-pyridin-3-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-butyramide;
- N*-{1-[6-(4-Chlorophenylsulfanyl)-pyridin-3-ylmethyl]-2,3-dihydro-1H-indol-5-yl}-3,3-dimethylbutyramide;
- N*-{1-[6-(4-Cyanophenoxy)-pyridin-3-ylmethyl]-2,3-dihydro-1H-indol-5-yl}-3,3-dimethylbutyramide;
- 10    3,3-Dimethyl-*N*-[1-(6-trifluoromethylpyridin-3-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-butyramide;
- 3,3-Dimethyl-*N*-[1-(3-methyl-benzo[*b*]thiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-butyramide;
- 15    *N*-[1-(6-Fluoro-4H-benzo[1,3]dioxin-8-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide;
- 3,3-Dimethyl-*N*-[1-(6-phenoxy-pyridin-3-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-butyramide;
- 3,3-Dimethyl-*N*-[1-(3-methyl-5-phenyl-isoxazol-4-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-butyramide;
- 20    *N*-(1-Benzo[*b*]thiophen-2-ylmethyl-2,3-dihydro-1H-indol-5-yl)-3,3-dimethylbutyramide;
- N*-{1-[1-(4-Fluorophenyl)-5-methyl-1H-pyrazol-4-ylmethyl]-2,3-dihydro-1H-indol-5-yl}-3,3-dimethylbutyramide;
- 25    3,3-Dimethyl-*N*-[1-(5-methylthiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-butyramide;
- 3,3-Dimethyl-*N*-[1-(4-pyrrol-1-yl-benzyl)-2,3-dihydro-1H-indol-5-yl]-butyramide;
- N*-[1-(4-Chlorobenzyl)-2,3-dihydro-1H-indol-5-yl]-2-(4-fluorophenyl)-acetamide;
- 2-(4-Fluorophenyl)-*N*-[1-(4-trifluoromethylbenzyl)-2,3-dihydro-1H-indol-5-yl]-acetamide;
- 30    2-(4-Fluorophenyl)-*N*-[1-(4-isopropylbenzyl)-2,3-dihydro-1H-indol-5-yl]-acetamide;
- 2-(4-Fluorophenyl)-*N*-[1-(3-fluoro-4-trifluoromethylbenzyl)-2,3-dihydro-1H-indol-5-yl]-acetamide;

- N*-[1-(6-Chlorobenzo[1,3]dioxol-5-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-2-(4-fluorophenyl)-acetamide;
- N*-[1-(3,5-Dimethyl-1-phenyl-1*H*-pyrazol-4-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-2-(4-fluorophenyl)-acetamide;
- 5 *N*-[1-(2-Chloro-5-trifluoromethylbenzyl)-2,3-dihydro-1*H*-indol-5-yl]-2-(4-fluorophenyl)-acetamide;
- N*-{1-[5-(4-Chlorophenoxy)-1,3-dimethyl-1*H*-pyrazol-4-ylmethyl]-2,3-dihydro-1*H*-indol-5-yl}-2-(4-fluorophenyl)-acetamide;
- N*-{1-[6-(4-Cyanophenoxy)-pyridin-3-ylmethyl]-2,3-dihydro-1*H*-indol-5-yl}-2-(4-fluorophenyl)-acetamide;
- 10 2-(4-Fluorophenyl)-*N*-[1-(3-methyl-benzo[*b*]thiophen-2-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-acetamide;
- N*-[1-(6-Fluoro-4*H*-benzo[1,3]dioxin-8-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-2-(4-fluorophenyl)-acetamide;
- 15 2-(4-Fluorophenyl)-*N*-[1-(6-phenoxy-pyridin-3-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-acetamide;
- N*-(1-Benzo[*b*]thiophen-2-ylmethyl-2,3-dihydro-1*H*-indol-5-yl)-2-(4-fluorophenyl)-acetamide;
- 2-(4-Fluorophenyl)-*N*-{1-[1-(4-fluorophenyl)-5-methyl-1*H*-pyrazol-4-ylmethyl]-2,3-dihydro-1*H*-indol-5-yl}-acetamide;
- 20 2-(4-Fluorophenyl)-*N*-[1-(5-methylthiophen-2-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-acetamide; and
- 2-(4-Fluorophenyl)-*N*-[1-(4-pyrrol-1-yl-benzyl)-2,3-dihydro-1*H*-indol-5-yl]-acetamide.

25

According to one embodiment, the invention relates to a pharmaceutical composition comprising one or more pharmaceutically acceptable carriers or diluents and a compound of formula I wherein the dotted line, s, q, U, X, Z, Y, R<sup>1</sup>, R<sup>1'</sup>, R<sup>2</sup> and R<sup>3</sup> are as defined above, accordingly any of the dotted line, a, b, c, d, e, f, g, h, j, k, s, q, L, T, U, X, Z, Y, W, R<sup>1</sup>, R<sup>1'</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>6'</sup>, R<sup>7</sup>, R<sup>7'</sup>, R<sup>8</sup>, R<sup>9</sup>, R<sup>9'</sup>, R<sup>10</sup>, R<sup>10'</sup>, R<sup>11</sup>, R<sup>12</sup> and R<sup>12'</sup> are as defined under formula I, or salts thereof. Pharmaceutical compositions of the invention may thus comprise one or more compounds of formula

30

I or salts thereof, such as one compound of formula I or a salt thereof; or two compounds of formula I or salts thereof; or three compounds of formula I or salts thereof.

- 5 The invention thus provides a pharmaceutical composition for oral or parenteral administration, said pharmaceutical composition comprising at least one compound of formula I or a salt thereof in a therapeutically effective amount together with one or more pharmaceutically acceptable carriers or diluents.
- 10 In one aspect, the compounds of the invention may be administered as the only therapeutically effective compound.

In another aspect the compounds of the invention may be administered as a part of a combination therapy, i.e. the compounds of the invention may be administered in  
15 combination with other therapeutically effective compounds having e.g. anti-convulsive properties. The effects of such other compounds having anti-convulsive properties may include but not be limited to activities on:

- ion channels such as sodium, potassium, or calcium channels
- the excitatory amino acid systems e.g. blockade or modulation of NMDA  
20 receptors
- the inhibitory neurotransmitter systems e.g. enhancement of GABA release, or blockade of GABA-uptake or
- membrane stabilisation effects.

Current anti-convulsive medications include, but are not limited to, tiagabine,  
25 carbamazepine, sodium valproate, lamotrigine, gabapentin, pregabalin, ethosuximide, levetiracetam, phenytoin, topiramate, zonisamide as well as members of the benzodiazepine and barbiturate class.

In one aspect, the compounds of the invention have been found to have effect on  
30 potassium channels of the KCNQ family, in particular the KCNQ2 subunit.

In one embodiment, the invention relates to the use of one or more compounds according to the invention in a method of treatment. The disorder or condition to be

prevented, treated or inhibited is responsive to an increased ion flow in a potassium channel such as the KCNQ family potassium ion channels. Such disorder or condition is preferably a disorder or condition of the central nervous system.

- 5 The compounds of the invention are considered useful for increasing ion flow in a voltage-dependent potassium channel in a mammal such as a human.

- The compounds of the invention are considered useful for the prevention, treatment or inhibition of a disorder or condition being responsive to an increased ion flow in a potassium channel such as the KCNQ family potassium ion channels. Such disorder or condition is preferably a disorder or condition of the central nervous system.
- 10

- The compounds of the invention are thus considered useful for preventing, treating or inhibiting disorders or diseases such as seizure disorders, neuropathic and migraine pain disorders, anxiety disorders and neurodegenerative disorders.
- 15

- Accordingly, the compounds of the invention are considered useful for the prevention, treatment or inhibition of disorders or conditions such as convulsions, epilepsy, anxiety disorders, neuropathic pain and neurodegenerative disorders.
- 20

According to one particular embodiment, the compounds of the invention are thus considered to be useful for preventing, treating or inhibiting seizure disorders such as convulsions, epilepsy and status epilepticus.

- 25 In one embodiment, the compounds of the invention are considered useful in the prevention, treatment and inhibition of convulsions.

- In another embodiment, the compounds of the invention are considered useful in the prevention, treatment and inhibition of epilepsy, epileptic syndromes and epileptic seizures.
- 30

In yet another embodiment, the compounds of the invention are considered useful in the prevention, treatment and inhibition of anxiety disorders such as anxiety and

conditions and diseases related to panic attack, agoraphobia, panic disorder with agoraphobia, panic disorder without agoraphobia, agoraphobia without history of panic disorder, specific phobia, social phobia and other specific phobias, obsessive-compulsive disorder, posttraumatic stress disorder, acute stress disorders, generalized  
5 anxiety disorder, anxiety disorder due to general medical condition, substance-induced anxiety disorder, separation anxiety disorder, adjustment disorders, performance anxiety, hypochondriacal disorders, anxiety disorder due to general medical condition and substance-induced anxiety disorder and anxiety disorder not otherwise specified.

10 In yet another embodiment, the compounds of the invention are considered useful in the prevention, treatment and inhibition of anxiety disorders such as anxiety, generalized anxiety disorder, panic anxiety, obsessive compulsive disorder, social phobia, performance anxiety, post-traumatic stress disorder, acute stress reaction, adjustment disorders, hypochondriacal disorders, separation anxiety disorder,  
15 agoraphobia, specific phobias, anxiety disorder due to general medical condition and substance-induced anxiety disorder.

In yet another embodiment, the compounds of the invention are also considered useful in the prevention, treatment and inhibition of neuropathic pain and migraine pain  
20 disorders such as allodynia, hyperalgesic pain, phantom pain, neuropathic pain related to diabetic neuropathy and neuropathic pain related to migraine.

In yet another embodiment,, the compounds of the invention are considered useful in the prevention, treatment and inhibition of neurodegenerative disorders such as  
25 Alzheimer's disease; Huntington's chorea; multiple sclerosis; amyotrophic lateral sclerosis; Creutzfeld-Jakob disease; Parkinson's disease; encephalopathies induced by AIDS or infection by rubella viruses, herpes viruses, borrelia and unknown pathogens; trauma-induced neurodegenerations; neuronal hyperexcitation states such as in medicament withdrawal or intoxication; and neurodegenerative diseases of the  
30 peripheral nervous system such as polyneuropathies and polyneuritides.

In yet another embodiment, the compounds of the invention are considered useful in the prevention, treatment and inhibition of neurodegenerative disorders such as



Alzheimer's disease; Huntington's chorea; multiple sclerosis; amyotrophic lateral sclerosis; Creutzfeld-Jakob disease; Parkinson's disease; encephalopathies induced by AIDS or infection by rubella viruses, herpes viruses, borrelia and unknown pathogens; and trauma-induced neurodegenerations.

5

In yet another embodiment, the compounds of the invention are considered useful in the prevention, treatment and inhibition of neuronal hyperexcitation states such as in medicament withdrawal or intoxication.

10 The invention provides compounds showing effect in one or more of the following tests:

- "Relative efflux through the KCNQ2 channel"

Which is a measure of the potency of the compound at the target channel

- "Maximum electroshock"

15 Which is a measure of seizures induced by non-specific CNS stimulation by electrical means

- "Pilocarpine induced seizures"

Seizures induced by pilocarpine are often difficult to treat with many existing antiseizure medications and so reflect a model of "drug resistant seizures"

20 • "Electrical seizure-threshold tests" and "Chemical seizure-threshold tests"

These models measure the threshold at which seizures are initiated, thus being models that detect whether compounds could delay seizure initiation.

- "Amygdala kindling"

25 Which is used as a measure of disease progression, as in normal animals the seizures in this model get more severe as the animal receives further stimulations.

According to one particular aspect of the invention, the compounds are KCNQ2 active with an  $EC_{50}$  of less than 15000nM such as less than 10000nM as measured by the  
30 test "Relative efflux through the KCNQ2 channel" which is described below.  
According to another particular aspect of the invention, the compounds are KCNQ2 active with an  $EC_{50}$  of less than 2000nM such as less than 1500nM as measured by the test "Relative efflux through the KCNQ2 channel" which is described below.

According to yet another particular aspect of the invention, the compounds are KCNQ2 active with an  $EC_{50}$  of less than 200nM such as less than 150nM as measured by the test "Relative efflux through the KCNQ2 channel" which is described below.

- 5 According to one particular aspect of the invention, the compounds have an  $ED_{50}$  of less than 15 mg/kg in the test "Maximum electroshock" which is described below. According to another particular aspect of the invention, the compounds have an  $ED_{50}$  of less than 5 mg/kg in the test "Maximum electroshock" which is described below.
- 10 According to one particular aspect of the invention, the compounds have an  $ED_{50}$  of less than 5 mg/kg in the "Electrical seizure -threshold test" and "Chemical seizure -threshold test" which is described below.

Some compounds have few or clinically insignificant side effects. Some of the  
15 compounds are thus tested in models of the unwanted sedative, hypothermic and ataxic actions of the compounds.

Some of the compounds have a large therapeutic index between anticonvulsant efficacy and side-effects such as impairment of locomotor activity or ataxic effects as  
20 measured by performance on a rotating rod. This means that the compounds will expectedly be well tolerated in patients permitting high doses to be used before side effects are seen. Thereby compliance with the therapy will expectedly be good and administration of high doses may be permitted making the treatment more efficacious in patients who would otherwise have side effects with other medications.

25

### Definitions

The term heteroatom refers to a nitrogen, oxygen or sulphur atom.

30 Halogen means fluoro, chloro, bromo or iodo.

The expressions  $C_{1-6}$ -alk(en/yn)yl and  $C_{1-6}$ -alk(an/en/yn)yl mean a  $C_{1-6}$ -alkyl,  $C_{2-6}$ -alkenyl or a  $C_{2-6}$ -alkynyl group. The term  $C_{1-6}$ -alkyl refers to a branched or un-

branched alkyl group having from one to six carbon atoms inclusive, including but not limited to methyl, ethyl, 1-propyl, 2-propyl, 1-butyl, 2-butyl, 2-methyl-2-propyl and 2-methyl-1-propyl. Similarly, C<sub>2-6</sub>-alkenyl and C<sub>2-6</sub>-alkynyl, respectively, designate such groups having from two to six carbon atoms, including one double bond and one  
5 triple bond respectively, including but not limited to ethenyl, propenyl, butenyl, ethynyl, propynyl and butynyl.

The expression C<sub>1-3</sub>-alk(en/yn)yl means a C<sub>1-3</sub>-alkyl, C<sub>2-3</sub>-alkenyl or a C<sub>2-3</sub>-alkynyl group. The term C<sub>1-3</sub>-alkyl refers to a branched or un-branched alkyl group having  
10 from one to three carbon atoms inclusive, including but not limited to methyl, ethyl, 1-propyl and 2-propyl. Similarly, C<sub>2-3</sub>-alkenyl and C<sub>2-3</sub>-alkynyl, respectively, designate such groups having from two to three carbon atoms, including one double bond and one triple bond respectively, including but not limited to ethenyl, 1-propenyl, 2-propenyl, 3-propenyl, ethynyl, 1-propynyl and 3-propynyl.

15 The expressions C<sub>3-8</sub>-cycloalk(en)yl and C<sub>3-8</sub>-cycloalk(an/en)yl mean a C<sub>3-8</sub>-cycloalkyl- or cycloalkenyl group. The term C<sub>3-8</sub>-cycloalkyl designates a monocyclic or bicyclic carbocycle having three to eight C-atoms, including but not limited to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, etc. The term C<sub>3-8</sub>-cycloalkenyl  
20 designates a monocyclic or bicyclic carbocycle having three to eight C-atoms and including one double bond.

The expressions C<sub>3-6</sub>-cycloalk(en)yl and C<sub>3-6</sub>-cycloalk(an/en)yl mean a C<sub>3-6</sub>-  
25 cycloalkyl- or cycloalkenyl group. The term C<sub>3-6</sub>-cycloalkyl designates a monocyclic or bicyclic carbocycle having three to six C-atoms, including but not limited to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, etc.

The term heterocycloalk(en)yl designates a monocyclic or bicyclic ring system  
30 wherein the ring is formed by 4 to 8 atoms selected from 2-7 carbonatoms and 1 or 2 heteroatoms selected from N, S, or O.

When two substituents together with a carbon atom to which they are attached form a 3-8 membered saturated or unsaturated ring which optionally contains 1 or 2 heteroatoms, then a monocyclic ring system is formed by 3 to 8 atoms selected from 1-8 carbonatoms and 0-2 heteroatoms selected from N, S, or O. Examples of such ring systems are cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl.

The term halo-C<sub>1-6</sub>-alk(en/yn)yl designates C<sub>1-6</sub>-alk(en/yn)yl being substituted with one or more halogen atoms, including but not limited to trifluoromethyl. Similarly, halo-C<sub>3-8</sub>-cycloalk(en)yl designates C<sub>3-8</sub>-cycloalk(en)yl being substituted with one or more halogen atoms and halo-heterocycloalk(en)yl designates heterocycloalk(en)yl being substituted with one or more halogen atoms.

The term NR<sup>12</sup>R<sup>12'</sup>-C<sub>1-6</sub>-alk(en/yn)yl designates C<sub>1-6</sub>-alk(en/yn)yl being substituted with NR<sup>12</sup>R<sup>12'</sup>. The term NR<sup>12</sup>R<sup>12'</sup>-C<sub>3-8</sub>-cycloalk(en)yl designates C<sub>3-8</sub>-cycloalk(en)yl being substituted with NR<sup>12</sup>R<sup>12'</sup>. The term NR<sup>12</sup>R<sup>12'</sup>-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl designates C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl being substituted with NR<sup>12</sup>R<sup>12'</sup>. When any of NR<sup>12</sup>R<sup>12'</sup>-C<sub>1-6</sub>-alk(en/yn)yl, NR<sup>12</sup>R<sup>12'</sup>-C<sub>3-8</sub>-cycloalk(en)yl and NR<sup>12</sup>R<sup>12'</sup>-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl is optionally substituted, then any of C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl is optionally substituted with one or more substituents independently being C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl or Ar.

As used herein, the term acyl refers to formyl, C<sub>1-6</sub>-alk(en/yn)ylcarbonyl, C<sub>3-8</sub>-cycloalk(en)ylcarbonyl, Ar-carbonyl, Ar-C<sub>1-6</sub>-alk(en/yn)ylcarbonyl or a C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl-carbonyl group, wherein C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl and Ar are as defined above.

When two substituents together with a nitrogen atom to which they are attached form a 4-8 membered saturated or unsaturated ring which optionally contains 1, 2 or 3 further heteroatoms, then a monocyclic ring system is formed by 4 to 8 atoms selected from the nitrogen atom, 1-7 carbonatoms and 0-3 further heteroatoms selected from N, S, or O. Examples of such ring systems are azetidine, beta-lactame, pyrrolidine,

piperidine, piperazine, morpholine, pyrrole, oxazolidine, thiazolidine, imidazolidine, azetidine, beta-lactame, tetrazole and pyrazole.

When two adjacent substituents together with the aromatic group to which they are  
 5 attached form a 4-8 membered ring, which optionally contains one or two  
 heteroatoms, then a ring system is formed by 4-8 atoms selected from 3-8  
 carbonatoms and 0-2 heteroatoms selected from N, S, or O. Such two adjacent  
 substituents may together form:

10  $-(CH_2)_n-CH_2-$ ,  $-CH=CH-(CH_2)_m-$ ,  $-CH_2-CH=CH-(CH_2)_p-$ ,  $-CH=CH-CH=CH-$ ,  
 $-(CH_2)_n-O-$ ,  $-O-(CH_2)_m-O-$ ,  $-CH_2-O-(CH_2)_p-O-$ ,  $-CH_2-O-CH_2-O-CH_2-$ ,  
 $-(CH_2)_n-S-$ ,  $-S-(CH_2)_m-S-$ ,  $-CH_2-S-(CH_2)_p-S-$ ,  $-CH_2-S-CH_2-S-CH_2-$ ,  
 $-(CH_2)_n-NH-$ ,  $-NH-(CH_2)_m-NH-$ ,  $-CH_2-NH-(CH_2)_p-NH-$ ,  $-CH=CH-NH-$ ,  
 $-O-(CH_2)_m-NH-$ ,  $-CH_2-O-(CH_2)_p-NH-$  or  $-O-(CH_2)_p-NH-CH_2-$ ,  $-S-(CH_2)_m-NH-$ ,  
 $N=CH-NH-$ ,  $-N=CH-O-$  or  $-N=CH-S-$ , wherein  $m$  is 1, 2 or 3,  $n$  is 2, 3 or 4 and  $p$   
 15 is 1 or 2.

The term Ar refers to optionally substituted aromatic systems of 5-10 carbon atoms,  
 wherein 0, 1, 2, 3 or 4 carbon atoms may be replaced by heteroatoms independently  
 selected from N, S, or O. Examples of such Ar groups are optionally substituted  
 20 phenyl, optionally substituted naphthyl, optionally substituted pyridine, optionally  
 substituted pyrrole, optionally substituted pyrimidine, optionally substituted  
 quinoline, optionally substituted indole, optionally substituted thiophene, optionally  
 substituted furan, optionally substituted thiazole and optionally substituted oxazole.  
 Ar may be substituted with one or more substituents independently being hydroxy,  
 25 halogen,  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl,  
 halo- $C_{1-6}$ -alk(en/yn)yl,  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -alk(en/yn)yl, acyl, nitro or  
 cyano,  $-CO-NH-C_{1-6}$ -alk(en/yn)yl,  $-CO-N(C_{1-6}$ -alk(en/yn)yl) $_2$ ,  $-NH_2$ ,  $-NH-C_{1-6}$ -  
 alk(en/yn)yl,  $-N(C_{1-6}$ -alk(en/yn)yl) $_2$ ,  $-S-C_{1-6}$ -alk(en/yn)yl,  $-SO_2-C_{1-6}$ -alk(en/yn)yl, -  
 $SO_2N(C_{1-6}$ -alk(en/yn)yl) $_2$  and  $-SO_2NH-C_{1-6}$ -alk(en/yn)yl; or two adjacent substituents  
 30 may together with the aromatic group to which they are attached form a 4-8  
 membered ring, which optionally contains one or two heteroatoms and which may be  
 saturated or unsaturated.

The terms C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl, C<sub>1-6</sub>-alk(en/yn)yl-heterocycloalk(en)yl, Ar, Ar-C<sub>1-6</sub>-alk(en/yn)yl, Ar-C<sub>3-8</sub>-cycloalk(en)yl, Ar-heterocycloalk(en)yl, Ar-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, Ar-C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl, Ar-C<sub>1-6</sub>-alk(en/yn)yl-heterocycloalk(en)yl, C<sub>1-6</sub>-alk(en/yn)oxy, C<sub>2-6</sub>-alkenyloxy, C<sub>2-6</sub>-alkynyloxy, C<sub>3-8</sub>-cycloalk(en)oxy, C<sub>1-6</sub>-alk(en/yn)oxy-C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)oxy-C<sub>1-6</sub>-alk(en/yn)yl, C<sub>1-6</sub>-alk(en/yn)oxy-C<sub>3-8</sub>-cycloalk(en)yl, C<sub>1-6</sub>-alk(en/yn)oxy-heterocycloalk(en)yl, Ar-oxy-C<sub>1-6</sub>-alk(en/yn)yl, Ar-C<sub>1-6</sub>-alk(en/yn)oxy-C<sub>1-6</sub>-alk(en/yn)yl, C<sub>1-6</sub>-alk(en/yn)ylcarbonyl, C<sub>3-8</sub>-alk(en/yn)ylcarbonyl, Ar-carbonyl, Ar-C<sub>1-6</sub>-alk(en/yn)ylcarbonyl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)ylcarbonyl, -CO-C<sub>1-6</sub>-alk(en/yn)yl, -S-C<sub>1-6</sub>-alk(en/yn)yl, -SO<sub>2</sub>-C<sub>1-6</sub>-alk(en/yn)yl and -SO<sub>2</sub>O-C<sub>1-6</sub>-alk(en/yn)yl, C<sub>1-6</sub>-alk(en/yn)oxy-carbonyl-C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)oxy-carbonyl-C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)oxy-carbonyl-C<sub>1-6</sub>-alk(en/yn)yl, acyl, acyl-C<sub>1-6</sub>-alk(en/yn)yl, acyl-C<sub>3-8</sub>-cycloalk(en)yl, acyl-heterocycloalk(en)yl, acyl-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, acyl-C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl, acyl-C<sub>1-6</sub>-alk(en/yn)yl-heterocycloalk(en)yl, hydroxy-C<sub>1-6</sub>-alk(en/yn)yl, hydroxy-C<sub>3-8</sub>-cycloalk(en)yl, hydroxy-heterocycloalk(en)yl, hydroxy-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, hydroxy-C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl, hydroxy-C<sub>1-6</sub>-alk(en/yn)yl-heterocycloalk(en)yl, halo-C<sub>1-6</sub>-alk(en/yn)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl, halo-heterocycloalk(en)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, halo-C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl, halo-C<sub>1-6</sub>-alk(en/yn)yl-heterocycloalk(en)yl, halo-C<sub>1-6</sub>-alk(en/yn)yl-Ar, halo-C<sub>3-8</sub>-cycloalk(en)yl-Ar, halo-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl-Ar, halo-C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl-Ar, halo-heterocycloalk(en)yl-Ar, cyano-C<sub>1-6</sub>-alk(en/yn)yl, cyano-C<sub>3-8</sub>-cycloalk(en)yl, cyano-heterocycloalk(en)yl, cyano-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, cyano-C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl, cyano-C<sub>1-6</sub>-alk(en/yn)yl-heterocycloalk(en)yl etc. designate such groups in which the C<sub>1-6</sub>-alk(en/yn)yl, C<sub>2-6</sub>-alkenyl, C<sub>2-6</sub>-alkynyl, C<sub>3-8</sub>-cycloalk(en)yl, heterocycloalk(en)yl, Ar, cyano, halo-C<sub>1-6</sub>-alk(en/yn)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl, halo-heterocycloalk(en)yl and acyl are as defined above.

The salts of the invention are preferably pharmaceutically acceptable salts. Such salts include pharmaceutical acceptable acid addition salts, pharmaceutically acceptable metal salts, ammonium and alkylated ammonium salts.

- 5 The pharmaceutically acceptable salts of the invention are preferably acid addition salts. The acid addition salts of the invention are preferably pharmaceutically acceptable salts of the compounds of the invention formed with non-toxic acids. Acid addition salts include salts of inorganic acids as well as organic acids. Representative examples of suitable inorganic acids include hydrochloric,
- 10 hydrobromic, hydroiodic, sulfuric, sulfamic, phosphoric and nitric acids and the like. Representative examples of suitable organic acids include formic, acetic, trichloroacetic, trifluoroacetic, propionic, benzoic, cinnamic, citric, fumaric, glycolic, lactic, maleic, malic, malonic, mandelic, oxalic, picric, pyruvic, salicylic, succinic, ethanesulfonic, tartaric, ascorbic, pantoic, gluconic, citraconic, aspartic, stearic,
- 15 palmitic, EDTA, glycolic, p-aminobenzoic, glutamic, bis-methylenesalicylic, methanesulfonic, ethanedisulfonic, itaconic, benzenesulfonic, p-toluenesulfonic acids, theophylline acetic acids, as well as the 8-halotheophyllines, for example 8-bromotheophylline and the like. Further examples of pharmaceutical acceptable inorganic or organic acid addition salts include the pharmaceutically acceptable salts
- 20 listed in J. Pharm. Sci. 1977,66,2, which is incorporated herein by reference.

Examples of metal salts include lithium, sodium, potassium, magnesium salts and the like.

- 25 Examples of ammonium and alkylated ammonium salts include ammonium, methyl-, dimethyl-, trimethyl-, ethyl-, hydroxyethyl-, diethyl-, n-butyl-, sec-butyl-, tert-butyl-, tetramethylammonium salts and the like.

- Also intended as pharmaceutically acceptable acid addition salts are the hydrates,
- 30 which the present compounds are able to form.

The compounds of the present invention may have one or more asymmetric centres and it is intended that any optical isomers, as separated, pure or partially purified

optical isomers or racemic mixtures thereof are included within the scope of the invention.

Furthermore, when a double bond or a fully or partially saturated ring system is present in the molecule geometric isomers may be formed. It is intended that any geometric isomers, as separated, pure or partially purified geometric isomers or mixtures thereof are included within the scope of the invention. Likewise, molecules having a bond with restricted rotation may form geometric isomers. These are also intended to be included within the scope of the present invention.

Furthermore, some of the compounds of the present invention may exist in different tautomeric forms and it is intended that any tautomeric forms that the compounds are able to form are included within the scope of the present invention.

The compounds of this invention may exist in unsolvated as well as in solvated forms with solvents such as water, ethanol and the like. In general, the solvated forms are considered equivalent to the unsolvated forms for the purposes of this invention. Racemic forms can be resolved into the optical antipodes by known methods, for example, by separation of diastereomeric salts thereof with an optically active acid, and liberating the optically active amine compound by treatment with a base. Another method for resolving racemates into the optical antipodes is based upon chromatography on an optically active matrix. Racemic compounds of the present invention can also be resolved into their optical antipodes, e.g. by fractional crystallization of d- or l- (tartrates, mandelates or camphorsulphonate) salts. The compounds of the present invention may also be resolved by the formation of diastereomeric derivatives.

Additional methods for the resolution of optical isomers, known to those skilled in the art, may be used. Such methods include those discussed by J. Jaques, A. Collet and S. Wilen in "Enantiomers, Racemates, and Resolutions", John Wiley and Sons, New York (1981).



Optically active compounds can also be prepared from optically active starting materials.

The invention also encompasses prodrugs of the present compounds, which on  
5 administration undergo chemical conversion by metabolic processes before becoming pharmacologically active substances. In general, such prodrugs will be functional derivatives of the compounds of the general formula I, VII, VIII, IX, X, XI, XXXII or XXXIII which are readily convertible in vivo into the required compound of the formula I, VII, VIII, IX, X, XI, XXXII or XXXIII. Conventional procedures for the  
10 selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs", ed. H. Bundgaard, Elsevier, 1985.

The invention also encompasses active metabolites of the present compounds.

15 Whenever mentioned in relation to the compounds of the formulas I, VII, VIII, IX, X, XI, XXXII or XXXIII, the terms epilepsy and epilepsies embrace any of the epilepsies, epileptic syndromes and epileptic seizures referred to in International League Against Epilepsy: Proposal for revised clinical and electroencephalographic classification of epileptic seizures. Commission on Classification and Terminology of  
20 the International League Against Epilepsy. *Epilepsia* 1981 22: 489-501 and in International League Against Epilepsy: Proposal for revised classification of epilepsies and epileptic syndromes. Commission on Classification and Terminology of the International League Against Epilepsy. *Epilepsia* 1989 30(4): 389-399.

25 Whenever mentioned in relation to the compounds of the formulas I, VII, VIII, IX, X; XI, XXXII or XXXIII, the term anxiety disorders embraces conditions and diseases related to panic attack, agoraphobia, panic disorder with agoraphobia, panic disorder without agoraphobia, agoraphobia without history of panic disorder, specific phobia, social phobia, obsessive-compulsive disorder, posttraumatic stress disorder,  
30 acute stress disorders, generalized anxiety disorder, anxiety disorder due to general medical condition, substance-induced anxiety disorder, separation anxiety disorder, adjustment disorders and anxiety disorder not otherwise specified as defined by

American Psychiatric Association *Diagnostic and statistical manual of mental disorders*, 4ed 1994: 110-113, 393-444 and 623-627.

### **Pharmaceutical compositions**

- 5 The compounds of this invention are generally utilized as the free base or as a pharmaceutically acceptable salt thereof. Representative examples are mentioned above.

- 10 If desired, the pharmaceutical composition of the invention may comprise the compound of formula I in combination with further pharmacologically active substances such as those described in the foregoing.

- The compounds of the invention may be administered alone or in combination with pharmaceutically acceptable carriers or excipients, in either single or multiple doses.
- 15 The pharmaceutical compositions according to the invention may be formulated with pharmaceutically acceptable carriers or diluents as well as any other known adjuvants and excipients in accordance with conventional techniques such as those disclosed in Remington: The Science and Practice of Pharmacy, 19 Edition, Gennaro, Ed., Mack Publishing Co., Easton, PA, 1995.

- 20 The pharmaceutical compositions may be specifically formulated for administration by any suitable route such as the oral, rectal, nasal, pulmonary, topical (including buccal and sublingual), transdermal, intracisternal, intraperitoneal, vaginal and parenteral (including subcutaneous, intramuscular, intrathecal, intravenous and
- 25 intradermal) route, the oral route being preferred. It will be appreciated that the preferred route will depend on the general condition and age of the subject to be treated, the nature of the condition to be treated and the active ingredient chosen.

- Pharmaceutical compositions for oral administration include solid dosage forms such
- 30 as capsules, tablets, dragees, pills, lozenges, powders and granules. Where appropriate, they can be prepared with coatings such as enteric coatings or they can be formulated so as to provide controlled release of the active ingredient such as sustained or prolonged release according to methods well known in the art.

Liquid dosage forms for oral administration include solutions, emulsions, suspensions, syrups and elixirs.

5    Pharmaceutical compositions for parenteral administration include sterile aqueous and nonaqueous injectable solutions, dispersions, suspensions or emulsions as well as sterile powders to be reconstituted in sterile injectable solutions or dispersions prior to use. Depot injectable formulations are also contemplated as being within the scope of the present invention.

10

Other suitable administration forms include suppositories, sprays, ointments, cremes, gels, inhalants, dermal patches, implants etc.

15

The pharmaceutical compositions of this invention or those which are manufactured in accordance with this invention may be administered by any suitable route, for example orally in the form of tablets, capsules, powders, syrups, etc., or parenterally in the form of solutions for injection. For preparing such compositions, methods well known in the art may be used, and any pharmaceutically acceptable carriers, diluents, excipients or other additives normally used in the art may be used.

20

25

A typical oral dosage is in the range of from about 0.001 to about 100 mg/kg body weight per day, preferably from about 0.01 to about 50 mg/kg body weight per day, and more preferred from about 0.05 to about 10 mg/kg body weight per day administered in one or more dosages such as 1 to 3 dosages. The exact dosage will depend upon the frequency and mode of administration, the sex, age, weight and general condition of the subject treated, the nature and severity of the condition treated and any concomitant diseases to be treated and other factors evident to those skilled in the art.

30

The formulations may conveniently be presented in unit dosage form by methods known to those skilled in the art. A typical unit dosage form for oral administration one or more times per day such as 1 to 3 times per day may contain from 0.05 to about 1000 mg, preferably from about 0.1 to about 500 mg, and more preferred from

about 0.5 mg to about 200 mg.

For parenteral routes such as intravenous, intrathecal, intramuscular and similar administration, typically doses are in the order of about half the dose employed for  
5 oral administration.

The compounds of this invention are generally utilized as the free substance or as a pharmaceutically acceptable salt thereof. One example is a base addition salt of a compound having the utility of a free acid. When a compound of the invention  
10 contains a free acid such salts may be prepared in a conventional manner by treating a solution or suspension of a free acid of the compound of the invention with a chemical equivalent of a pharmaceutically acceptable base. Representative examples are mentioned above.

15 For parenteral administration, solutions of the novel compounds of the invention in sterile aqueous solution, aqueous propylene glycol, aqueous vitamin E or sesame or peanut oil may be employed. Such aqueous solutions should be suitably buffered if necessary and the liquid diluent first rendered isotonic with sufficient saline or glucose. The aqueous solutions are particularly suitable for intravenous,  
20 intramuscular, subcutaneous and intraperitoneal administration. The sterile aqueous media employed are all readily available by standard techniques known to those skilled in the art.

Solutions for injections may be prepared by dissolving the active ingredient and  
25 possible additives in a part of the solvent for injection, preferably sterile water, adjusting the solution to a desired volume, sterilising the solution and filling it in suitable ampules or vials. Any suitable additive conventionally used in the art may be added, such as tonicity agents, preservatives, antioxidants, etc.

30 Suitable pharmaceutical carriers include inert solid diluents or fillers, sterile aqueous solution and various organic solvents.

Examples of solid carriers are lactose, terra alba, sucrose, cyclodextrin, talc, agar, pectin, acacia, stearic acid and lower alkyl ethers of cellulose corn starch, potato starch, talcum, magnesium stearate, gelatine, lactose, gums, and the like.

- 5 Any other adjuvants or additives usually used for such purposes such as colourings, flavourings, preservatives etc. may be used provided that they are compatible with the active ingredients.

- 10 Examples of liquid carriers are syrup, peanut oil, olive oil, phospho lipids, fatty acids, fatty acid amines, polyoxyethylene and water. Similarly, the carrier or diluent may include any sustained release material known in the art, such as glyceryl monostearate or glyceryl distearate, alone or mixed with a wax.

- 15 The pharmaceutical compositions formed by combining the novel compounds of the invention and the pharmaceutical acceptable carriers are then readily administered in a variety of dosage forms suitable for the disclosed routes of administration. The formulations may conveniently be presented in unit dosage form by methods known in the art of pharmacy.

- 20 Formulations of the present invention suitable for oral administration may be presented as discrete units such as capsules or tablets, each containing a predetermined amount of the active ingredient, and which may include one or more suitable excipients. Furthermore, the orally available formulations may be in the form of a powder or granules, a solution or suspension in an aqueous or non-aqueous  
25 liquid, or an oil-in-water or water-in-oil liquid emulsion.

If a solid carrier is used for oral administration, the preparation may be tablette, placed in a hard gelatine capsule in powder or pellet form or it can be in the form of a troche or lozenge.

30

The amount of solid carrier will vary widely but will usually be from about 25 mg to about 1 g.

If a liquid carrier is used, the preparation may be in the form of a syrup, emulsion, soft gelatine capsule or sterile injectable liquid such as an aqueous or non-aqueous liquid suspension or solution.

- 5 If desired, the pharmaceutical composition of the invention may comprise the compound of the formula **I, VII, VIII, IX, X or XI** in combination with further pharmacologically active substances such as those described in the foregoing.

Typical examples of recipes for the formulation of the invention are as follows:

10

- 1) Tablets containing 5.0 mg of a compound of the invention calculated as the free base:

Compound of formula <b>I, VII, VIII, IX, X or XI</b>	5.0 mg
Lactose	60 mg
Maize starch	30 mg
Hydroxypropylcellulose	2.4 mg
Microcrystalline cellulose	19.2 mg
Croscarmellose Sodium Type A	2.4 mg
Magnesium stearate	0.84 mg

20

- 2) Tablets containing 0.5 mg of a compound of the invention calculated as the free base:

Compound of formula <b>I, I, VII, VIII, IX, X or XI</b>	0.5 mg
Lactose	46.9 mg
Maize starch	23.5 mg
Povidone	1.8 mg
Microcrystalline cellulose	14.4 mg
Croscarmellose Sodium Type A	1.8 mg
Magnesium stearate	0.63 mg

30


- 3) Syrup containing per millilitre:

Compound of formula <b>I, VII, VIII, IX, X or XI</b>	25 mg
Sorbitol	500 mg

		Hydroxypropylcellulose	15 mg	
		Glycerol	50 mg	
		Methyl-paraben	1 mg	
		Propyl-paraben	0.1 mg	
5		Ethanol	0.005 mL	
		Flavour	0.05 mg	
		Saccharin sodium	0.5 mg	
		Water	ad 1 mL	
10	4)	Solution for injection containing per millilitre:		
		Compound of formula I, VII, VIII, IX, X or XI		0.5 mg
		Sorbitol	5.1 mg	
		Acetic Acid	0.05 mg	
		Saccharin sodium	0.5 mg	
15		Water		


## Preparation of the compounds of the invention

Scheme 1



5

Scheme 2





The compounds of the invention of the general formula **I**, wherein the the dotted line, **q**, **s**, **U**, **Y**, **X**, **Z**, **R<sup>1</sup>**, **R<sup>1'</sup>**, **R<sup>2</sup>** and **R<sup>3</sup>** are as defined above, accordingly any of dotted line, **a**, **b**, **c**, **d**, **e**, **f**, **g**, **h**, **s**, **q**, **U**, **X**, **Z**, **Y**, **W**, **R<sup>1</sup>**, **R<sup>1'</sup>**, **R<sup>2</sup>**, **R<sup>3</sup>**, **R<sup>5</sup>**, **R<sup>6</sup>**, **R<sup>6'</sup>**, **R<sup>7</sup>**, **R<sup>7'</sup>**, **R<sup>8</sup>**, **R<sup>9</sup>**, **R<sup>9'</sup>**, **R<sup>10</sup>**, **R<sup>10'</sup>**, **R<sup>11</sup>**, **R<sup>12</sup>** and **R<sup>12'</sup>** are defined under formula **I** are prepared by the methods as described below and as represented in the Schemes 1 and 2.

Indoles and indolines of the general formula **XII** and **XIII** substituted at position 4 or 6 with **R<sup>2</sup>-(U)<sub>s</sub>** are commercially available, described in the literature or prepared according to methods known to chemists skilled in the art [R. J. Sundberg "Pyrroles and their Benzo Derivatives: (iii) Synthesis and Applications" in *Comprehensive Heterocyclic Chemistry*, A. R. Katritzky, C.W. Rees (Editors), vol. IV, pp 313-376, Pergamon Press, 1984]. Indoles of the general formula **XII** can be converted into indolines of the general formula **XIII** by methods known to chemists skilled in the art such as catalytic hydrogenation or reduction with NaBH<sub>3</sub>CN in appropriate solvents such as acetic acid [S. M. Bromidge, S. Dabbs, D. T. Davies, D. M. Duckworth, I. T. Forbes et al. *J. Med. Chem.* **41**, 1998, 1598-1612]. Compounds of the general formula **XII** or **XIII** with **s** being 0 and **R<sup>2</sup>** being in particular but not limited to substituted aryl or substituted heteroaryl as defined above can be prepared from corresponding compounds with **R<sup>2</sup>** being I or Br by means of C-C coupling reactions known to chemists skilled in the art, such as *Suzuki* coupling, *Stille* coupling, or other transition metal catalysed cross-coupling reactions [D.W. Knight "Coupling Reactions Between sp<sup>2</sup> Carbon Centers" in *Comprehensive Organic Synthesis*, v. 3, pp. 481-520, Pergamon Press 1991].

Compounds of the general formula **XIV** are prepared by protection of the indoline nitrogen of the compounds of the general formula **XIII** with an appropriate protecting group (PG<sup>1</sup>) [*Protective Groups in Organic Synthesis*, 3rd Edition T. W. Greene, P. G. M. Wuts, Wiley Interscience 1999], such as a trifluoroacetyl group known to chemists skilled in the art as TFA group, by reaction with the reagent forming the protective group such as trifluoroacetic acid anhydride in a suitable solvent, such as 1,2-dichloroethane at appropriate temperatures.

The obtained compounds of the general formula XIV are converted into compounds of the general formula XV by regioselective nitration at position 5 of the indoline moiety by methods known to chemists skilled in the art [R. Behnisch "Aromatische Nitro-Verbindungen" in *Methoden der Organische Chemie/(Houben-Weyl)* p. 255, v. E16d, Thieme: 1992] such as reaction with concentrated nitric acid in appropriate solvent such as acetic anhydride, acetic acid, concentrated sulphuric acid or mixtures thereof at appropriate temperatures. The nitro compounds of the general formula XV where  $R^2$  is halogen, in particular fluorine, and s is 0 can be converted into compounds of the general formula XV, where U is O,  $NR^{11}$  or S and  $R^2$  is as defined above, by nucleophilic aromatic substitution reactions known to chemists skilled in the art such as reaction with the appropriate nucleophiles forming the  $-(U)_s-R^2$  group such as thiophenols, alkylsulfides, alcohols, phenols, amines, and anilines in their neutral or deprotonated form. The compounds of the general formula XV where U is  $SO_2$  can be obtained from the compounds of the general formula XV, where U is S, by oxidation according to methods known to the chemist skilled in the art, for example by oxidation with  $NaIO_4$  in the presence of  $RuCl_3$  as a catalyst or with 3-chloroperoxybenzoic acid.

The nitro group in compounds of the general formula XV can be reduced with suitable reducing agents such as zinc or iron powder in the presence of acid such as acetic acid or aqueous hydrochloric acid, or hydrogen gas or ammonium formate in the presence of a suitable hydrogenation catalyst such as palladium on activated carbon in suitable solvents such as methanol, ethanol, or tetrahydrofuran, at suitable temperatures or under ultrasonic irradiation, to obtain anilines of the general formula XVI. Alternatively, tin (II) chloride or sodium dithionite can be used as reducing agents under conditions well known to the chemist skilled in the art.

Compounds of the general formula XVII are prepared from compounds of the general formula XVI by the reaction with suitable electrophilic reagents forming an  $R^3-(Z)_q-X$  group, such as alkyl, aryl or heteroaryl chloroformates or carbamyl chlorides, acid chlorides, acid bromides, acid iodides, sulfonyl chlorides, isocyanates, carbonic acid anhydrides, activated carbonic acids with activating reagents such as carbodiimides or others as known to chemists skilled in the art in suitable solvents, such as acetonitrile,

tetrahydrofuran, 1,2-dichloroethane, or methylene chloride, at suitable temperature, such as room temperature or reflux, with or without addition of bases, such as magnesium oxide, potassium carbonate, sodium hydride, trialkylamines, sodium- or potassium alcoholates, or pyridine, reactions well known to the chemist skilled in the art. Then the protective group  $PG^1$  is removed according to methods known to chemists skilled in the art [*Protective Groups in Organic Synthesis*, 3rd Edition T. W. Greene, P. G. M. Wuts, Wiley Interscience 1999], furnishing compounds of the general formula XVIII. For example, when  $PG^1$  is TFA, it can be removed by hydrolysis with aqueous potassium carbonate in an appropriate solvent, such as methanol, at a suitable temperature.

Finally, the obtained anilines of the general formula XVIII are subjected to reductive alkylation reactions, known to chemists skilled in the art, with aldehydes of the general formula  $YCHO$  where  $Y$  is defined as above in the presence of suitable reducing agent such as  $NaBH_3CN$  in suitable solvents such as methanol, ethanol, tetrahydrofuran, acetonitrile or mixtures thereof, with or without addition of catalytic amounts of acid, such as acetic acid, at suitable temperatures forming compounds of the invention of the general formula I, where  $R^1$  and  $R^{1'}$  are hydrogens. Alternatively, a  $(Y)(R^1)(R^{1'})C-$  group can be introduced by nucleophilic substitution reactions with the appropriate electrophiles of the general formula  $(Y)(R^1)(R^{1'})C-LG$ , where  $LG$  is a suitable leaving group such as iodide, bromide, or sulphonate, under conditions known to the chemist skilled in the art, furnishing the compounds of the invention of the general formula I.

Alternatively, compounds of the general formula XIX are commercially available, described in the literature or can be prepared from compounds of the general formula XV by deprotection as described above. Then they are subjected to reductive alkylation with aldehydes of the general formula  $YCHO$  or to nucleophilic substitution reactions with electrophiles of the general formula  $(Y)(R^1)(R^{1'})C-LG$  as described above, furnishing compounds of the general formula XX. Then the nitro group is reduced as described above forming compounds of the general formula XXI. Finally, the compounds of the invention of the general formula I with indoline moiety

are obtained by the method described above for the conversion of compounds of the general formula **XVI** into compounds of the general formula **XVII**.

Optionally, compounds of the invention of the general formula **I** with indole moiety can be obtained from indolines of the general formula **I** by means of dehydrogenation known to chemists skilled in the art such as oxidation with appropriate reagents such as 2,3,5,6-tetrachloro-[1,4]benzoquinone,  $\text{MnO}_2$ , or catalytic dehydrogenation in the presence of a catalyst such as Pd on charcoal or  $\text{RuCl}_2(\text{PPh}_3)_3$  in appropriate solvents such as toluene or xylene at appropriate temperatures.

Alternatively, compounds of the general formula **I** where  $-(\text{U})_s-\text{R}^2$  is attached to the position 6 of the indoline moiety, can be prepared by a route shown in Scheme 2 as follows:

5-Nitroindoline is protected with an appropriate protecting group, such as TFA group, as described above for compounds of the general formula **XIV**, furnishing compounds of the general formula **XXII**. Then the nitro group is reduced as described above for preparation of compounds of the general formula **XVI**, furnishing compounds of the general formula **XXIII**. They are converted into compounds of the general formula **XXIV** with appropriate electrophiles forming  $\text{R}^3-(\text{Z})_q-\text{X}$  as described above for compounds of the general formula **XVII**. Compounds of the general formula **XXV** where  $s$  is 0 and  $\text{R}^2$  is  $\text{NO}_2$  or halogen such as Cl, Br or I, are obtained by means of regioselective electrophilic aromatic substitution, well known to chemists skilled in the art, with appropriate electrophiles such as N-chlorosuccinimide, bromine, iodine, iodochloride in the appropriate solvent such as acetic acid or by nitration under conditions as described for compounds of the general formula **XV**.

Compounds of the general formula **XXV** where  $s$  is 0 and  $\text{R}^2$  is substituted aryl or substituted heteroaryl as defined above can be prepared from corresponding compounds of the same general formula where  $\text{R}^2$  is I or Br by means of C-C coupling reactions known to chemists skilled in the art as described above. Then the protective group is removed as described above, furnishing the compounds of the general formula **XXVI**.

Finally, the compounds of the invention of the general formula I with indoline moiety are prepared from the compounds of the general formula XXVI by reductive alkylation or by nucleophilic substitution reactions as described above. Also, the compounds of the invention of the general formula I with indole moiety can be obtained from indolines of the general formula I by means of dehydrogenation as described above.

### Examples

Analytical LC-MS data were obtained on a PE Sciex API 150EX instrument equipped with an APPI (atmospheric pressure photo ionisation) ion source and Shimadzu LC-8A/SLC-10A LC system. Column: 30 X 4.6 mm Waters Symmetry C18 column with 3.5 µm particle size; Solvent system: A = water/trifluoroacetic acid (100:0.05) and B = water/acetonitrile/trifluoroacetic acid (5:95:0.03); Method: Linear gradient elution with 90% A to 100% B in 4 minutes and with a flow rate of 2 mL/minute. Purity was determined by integration of the UV (254 nm) and ELSD trace. The retention times (RT) are expressed in minutes.

Preparative LC-MS-purification was performed on the same instrument. Column: 50 X 20 mm YMC ODS-A with 5 µm particle size; Method: Linear gradient elution with 80% A to 100% B in 7 minutes and with a flow rate of 22.7 mL/minute. Fraction collection was performed by split-flow MS detection.

<sup>1</sup>H NMR spectra were recorded at 500.13 MHz on a Bruker Avance DRX500 instrument. Deuterated chloroform (99.8%D) or dimethyl sulfoxide (99.8%D) were used as solvents. TMS was used as internal reference standard. Chemical shift values are expressed in ppm-values. The following abbreviations are used for multiplicity of NMR signals: s = singlet, d = doublet, t = triplet, q = quartet, qui = quintet, h = heptet, dd = double doublet, dt = double triplet, dq = double quartet, tt = triplet of triplets, m = multiplet and br. = broad.

### Preparation of intermediates

#### Preparation of intermediates of the general formula XXII and XIV

*1-Trifluoroacetyl-5-nitroindoline.*

To a suspension of 5-nitroindoline (5.51 g, 33.56 mmol) in 1,2-dichloroethane (15 ml) trifluoroacetic anhydride (20 ml) was added. After 60 min the obtained solution was quenched with heptane (200 ml) and the title compound was separated by filtration in two crops. Yield 7.12 g, 81.5%. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 3.34 (t, 2H), 4.38 (t, 2H), 8.19 (m, 3H).

*1-Trifluoroacetyl-4-chloroindoline* was prepared analogously from 4-chloroindoline [S. M. Bromidge, S. Dabbs, D. T. Davies, D. M. Duckworth, I. T. Forbes et al. *J. Med. Chem.* **41**, 1998, 1598-1612]. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 3.26 (t, 2H), 4.34 (t, 2H), 7.27 (d, 1H), 7.34 (t, 1H), 8.01 (d, 1H).

**Preparation of intermediates of the general formula XV***1-Trifluoroacetyl-4-chloro-5-nitroindoline.*

To a solution of 1-trifluoroacetyl-4-chloroindoline (197 mg, 0.838 mmol) in acetic anhydride (3 ml) and acetic acid (0.3 ml) a solution of fuming HNO<sub>3</sub> (0.4 ml) was added by small portions during 5 hours. The resulting reaction mixture was poured into ice, neutralised with saturated aqueous NaHCO<sub>3</sub>, and extracted with ethyl acetate. The organic solution was filtered via plug of SiO<sub>2</sub> (10 g), evaporated in vacuo and purified by flash chromatography on SiO<sub>2</sub> with gradient heptane – 1:4 ethyl acetate/heptane to give 70 mg of the title compound as yellow solid, yield 31%. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 3.33 (t, 2H), 4.42 (t, 2H), 8.10 (s, 2H).

**Preparation of intermediates of the general formula XX***1-(5-Chlorothiophen-2-ylmethyl)-5-nitroindoline.*

To a solution of 5-nitroindoline (3.23 g, 19.67 mmol) and 5-chlorothiophene-2-carboxaldehyde (4.2 g, 28.6 mmol) in methanol (45 ml) and acetic acid (8 ml) a solution of NaBH<sub>3</sub>CN (0.9 g) in methanol (8 ml) was added dropwise during 10 min. The obtained reaction mixture was stirred overnight. The title compound was

separated by filtration, washed with methanol and water and dried in vacuo to furnish 4.6 g of red crystalline solid. Yield 79.3%. LC/MS (m/z) 293.9 ( $[M]^+$ ); RT = 3.59, (UV, ELSD) 98%, 99.8%.  $^1\text{H}$  NMR (DMSO- $d_6$ ): 3.04 (t, 2H), 3.62 (t, 2H), 4.68 (s, 2H), 6.72 (d, 1H), 6.98 (d, 1H), 7.01 (d, 1H), 7.85 (unresolved m, 1H), 8.00 (dd, 1H).

5

The following compound was prepared analogously using appropriate aldehydes:

*1-(4-Fluorobenzyl)-5-nitroindoline.*

Yellow needles, yield 3.66 g, 72.2%. LC/MS (m/z) 272.0 ( $[M]^+$ ); RT = 3.35, (UV, ELSD) 99%, 100%.  $^1\text{H}$  NMR (DMSO- $d_6$ ): 3.06 (t, 2H), 3.61 (t, 2H), 4.52 (s, 2H), 6.63 (d, 1H), 7.18 (m, 2H), 7.35 (m, 2H), 7.83 (unresolved m, 1H), 7.97 (dd, 1H).

10

**Preparation of intermediates of the general formula XXI, XXII, and XVI**

15 *1-(5-Chlorothiophen-2-ylmethyl)-5-aminoindoline.*

To a cold (ice/water bath) vigorously stirred solution of 1-(5-Chlorothiophen-2-ylmethyl)-5-nitroindoline (4.013 g, 13.62 mmol) in THF (100 ml) and acetic acid (15 ml) zinc powder (25 g) was added by small portions maintaining the temperature below 40°C. The cold bath removed and the stirring continued at room temperature until reaction completion (1 hour). The obtained suspension was filtered via a plug of  $\text{SiO}_2$  (25 g) with ethyl acetate as an eluent and obtained solution was evaporated in vacuo. The obtained residue was treated with saturated aqueous  $\text{NaHCO}_3$ , extracted with ethyl acetate, dried over  $\text{Na}_2\text{SO}_4$  and evaporated in vacuo to give the title compound as a dark green oil. Yield 3.30 g, 91.5%. LC/MS (m/z) 265.9 ( $[M+1]^+$ ); RT = 1.85, (UV, ELSD) 93%, 100%.  $^1\text{H}$  NMR (DMSO- $d_6$ ): 2.73 (t, 2H), 3.08 (t, 2H), 4.25 (s, 2H), 4.40 (br. s, 2H,  $\text{NH}_2$ ), 6.30 (dd, 1H), 6.41 (d, 1H), 6.43 (unresolved m, 1H), 6.89 (d, 1H), 6.95 (d, 1H).

20

25

The following compounds were prepared analogously:

30

*1-(4-Fluorobenzyl)-5-aminoindoline.*

The obtained crude product after filtration via  $\text{SiO}_2$  was dissolved in a small amount of methanol, quenched with saturated aqueous  $\text{NaHCO}_3$ , and the title compound was

separated by filtration, washed with water, and dried in vacuo. Yield 2.40 g, 93.2%, dark violet solid. LC/MS (m/z) 265.9 ( $[M+1]^+$ ); RT = 1.74, (UV, ELSD) 87%, 98%.  $^1\text{H}$  NMR (DMSO- $d_6$ ): 2.72 (t, 2H), 3.01 (t, 2H), 4.04 (s, 2H), 4.36 (br. s, 2H,  $\text{NH}_2$ ), 6.28 (d, 1H), 6.34 (d, 1H), 6.44 (s, 1H), 7.14 (t, 2H), 7.38 (t, 2H).

5

*1-Trifluoroacetyl-5-aminoindoline.*

The title compound was prepared from 1-trifluoroacetyl-5-nitroindoline (6.67 g, 25.65 mmol). The crude product after filtration via  $\text{SiO}_2$  was used in the next step without purification. Yield 6.11 g, 100%. LC/MS (m/z) 230.1 ( $[M]^+$ ); RT = 1.29, (UV, ELSD) 97%, 98%.  $^1\text{H}$  NMR (DMSO- $d_6$ ): 3.10 (t, 2H), 4.18 (t, 2H), 5.18 (br. s, 2H,  $\text{NH}_2$ ), 6.43 (dd, 1H), 6.53 (s, 1H), 7.75 (d, 1H).

10

*1-Trifluoroacetyl-4-chloro-5-aminoindoline.*

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ): 3.23 (t, 2H), 4.28 (t, 2H), 6.67 (d, 1H), 7.93 (d, 1H).

15

**Preparation of intermediates of the general formula XXIV and XVII**

*3,3-Dimethyl-N-[1-(2,2,2-trifluoroacetyl)-2,3-dihydro-1H-indol-5-yl]-butyramide.*

To a cold (ice/water bath) solution of 1-trifluoroacetyl-5-aminoindoline (2.69 g, 11.7 mmol) in  $\text{CH}_2\text{Cl}_2$  *tert*-butylacetyl chloride (1.88 g, 14 mmol) was added followed by addition of  $\text{Et}_3\text{N}$  (4 ml). After 5 min the reaction mixture was quenched with saturated aqueous  $\text{NaHCO}_3$  and stirred for 30 min. The organic layer was filtered via plug of  $\text{SiO}_2$  (20 g) with ethyl acetate as an eluent and evaporated to a small volume. It was quenched with heptane and the title compound was separated by filtration. Yield 3.10 g, 81%, white solid. LC/MS (m/z) 329.2 ( $[M+1]^+$ ); RT = 3.04, (UV, ELSD) 97%, 100%.  $^1\text{H}$  NMR (DMSO- $d_6$ ): 1.02 (s, 9H), 2.18 (s, 2H), 3.22 (t, 2H), 4.26 (t, 2H), 7.38 (dd, 1H), 7.72 (s, 1H), 7.96 (d, 1H), 9.86 (s, 1H,  $\text{NHCO}$ ).

20

25

30

The following compounds were prepared analogously from 1-trifluoroacetyl-5-aminoindoline and appropriate acid chloride or chloroformate:

*N-[4-Chloro-1-(2,2,2-trifluoroacetyl)-2,3-dihydro-1H-indol-5-yl]-3,3-*

*dimethylbutyramide* was prepared from 1-trifluoroacetyl-4-chloro-5-aminoindoline.



The reaction mixture was evaporated and used in the next step without characterisation.

*2,2-Dimethyl-N-[1-(2,2,2-trifluoroacetyl)-2,3-dihydro-1H-indol-5-yl]-propionamide.*

- 5 <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 1.22 (s, 9H), 3.23 (t, 2H), 4.28 (t, 2H), 7.47 (dd, 1H), 7.71 (s, 1H), 7.96 (d, 1H), 9.26 (s, 1H, NHCO).

*2-(4-Fluorophenyl)-N-[1-(2,2,2-trifluoroacetyl)-2,3-dihydro-1H-indol-5-yl]-acetamide.*

- 10 LC/MS (m/z) 367.0 ([M+1]<sup>+</sup>); RT = 3.00, (UV, ELSD) 92%, 99%. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 3.22 (t, 2H), 3.63 (s, 2H), 4.27 (t, 2H), 7.15 (t, 2H), 7.36 (dd, 2H), 7.39 (dd, 1H), 7.69 (s, 1H), 7.97 (d, 1H), 10.24 (s, 1H, NHCO).

*[1-(2,2,2-Trifluoroacetyl)-2,3-dihydro-1H-indol-5-yl]-carbamic acid ethyl ester.*

- 15 The title compound was prepared using 1,2-dichloroethane as a solvent and pyridine as a base. LC/MS (m/z) 302.1 ([M]<sup>+</sup>); RT = 2.85 (UV, ELSD) 79%, 100%. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 1.24 (t, 3H), 3.22 (t, 2H), 4.12 (q, 2H), 4.26 (t, 2H), 7.31 (br. d (unresolved dd), 1H), 7.49 (s, 1H), 7.94 (d, 1H), 9.70 (s, 1H, NHCO).

- 20 *[1-(2,2,2-Trifluoroacetyl)-2,3-dihydro-1H-indol-5-yl]-carbamic acid propyl ester.*

The title compound was prepared using 1,2-dichloroethane as a solvent and pyridine as a base. LC/MS (m/z) 315.9 ([M]<sup>+</sup>); RT = 3.11 (UV, ELSD) 89%, 99%. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 0.93 (t, 3H), 1.64 (m, 2H), 3.22 (t, 2H), 4.03 (t, 2H), 4.26 (t, 2H), 7.32 (br. d (unresolved dd), 1H), 7.50 (s, 1H), 7.94 (d, 1H), 9.71 (s, 1H, NHCO).

25

#### **Preparation of intermediates of the general formula XXV and XXVI**

*3,3-Dimethyl-N-(6-nitro-2,3-dihydro-1H-indol-5-yl)-butyramide.*

- 30 To a cold (ice/water bath) stirred solution of 3,3-dimethyl-N-[1-(2,2,2-trifluoroacetyl)-2,3-dihydro-1H-indol-5-yl]-butyramide (1.96 g, 5.98 mmol) in acetic anhydride (30 ml) and acetic acid (5 ml) a solution of fuming HNO<sub>3</sub> (650 mg, 10.3 mmol) in acetic acid (5 ml) was added dropwise during 5 min. After 5 min the reaction mixture was poured into ice and neutralised with solid NaHCO<sub>3</sub> which was

added by small portions with stirring until gas formation ceased. The yellow solid of 3,3-dimethyl-N-[6-nitro-1-(2,2,2-trifluoroacetyl)-2,3-dihydro-1H-indol-5-yl]-butyramide was filtered, washed with water and dried in vacuo. LC/MS (m/z) 374.0 ( $[M+1]^+$ ); RT = 3.45 (UV, ELSD) 94%, 99%.  $^1\text{H}$  NMR (DMSO- $d_6$ ): 1.03 (s, 9H), 2.23 (s, 2H), 3.35 (t, 2H), 4.36 (t, 2H), 7.66 (s, 1H), 7.51 (s, 1H), 10.17 (s, 1H, NHCO).

The solid was redissolved in methanol (30 ml) followed by addition of  $\text{K}_2\text{CO}_3$  (2.0 g) in water (7 ml). The colour changed immediately from yellow to dark red. After stirring for 15 min the reaction mixture was poured into ice/water and the title compound was isolated by filtration to give 1.52 g of purple solid, yield 91.8%. LC/MS (m/z) 277.0 ( $[M]^+$ ); RT = 2.30 (UV, ELSD) 91%, 99%.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ): 1.10 (s, 9H), 2.29 (s, 2H), 3.10 (t, 2H), 3.63 (t, 2H), 4.80 (very br. s, NH), 7.30 (s, 1H), 8.46 (s, 1H), 10.14 (s, 1H, NHCO).

The following compounds were prepared analogously:

*2,2-Dimethyl-N-(6-nitro-2,3-dihydro-1H-indol-5-yl)-propionamide.*

LC/MS (m/z) 264.1 ( $[M+1]^+$ ); RT = 2.19 (UV, ELSD) 96%, 95%.  $^1\text{H}$  NMR (DMSO- $d_6$ ): 1.19 (s, 9H), 3.00 (t, 2H), 3.51 (dt, 2H), 5.98 (br. s, NH), 6.97 (s, 1H), 7.44 (s, 1H), 9.57 (s, 1H, NHCO).

*2-(4-Fluorophenyl)-N-(6-nitro-2,3-dihydro-1H-indol-5-yl)-acetamide.*

LC/MS (m/z) 315.0 ( $[M]^+$ ); RT = 2.33 (UV, ELSD) 87%, 99%.  $^1\text{H}$  NMR (DMSO- $d_6$ ): 2.99 (t, 2H), 3.49 (dt, 2H), 3.62 (s, 2H), 6.00 (br. s, NH), 6.91 (s, 1H), 7.15 (t, 2H), 7.28 (s, 1H), 7.33 (dd, 2H), 9.95 (s, 1H, NHCO).

*[6-Nitro-2,3-dihydro-1H-indol-5-yl]-carbamic acid ethyl ester.*

LC/MS (m/z) 250.9 ( $[M]^+$ ); RT = 1.92 (UV, ELSD) 93%, 98%.  $^1\text{H}$  NMR (DMSO- $d_6$ ): 1.19 (t, 3H), 2.99 (t, 2H), 3.50 (dt, 2H), 4.06 (q, 2H), 5.96 (br. s, NH), 6.92 (s, 1H), 7.24 (s, 1H), 9.22 (s, 1H, NHCO).

*[6-Nitro-2,3-dihydro-1H-indol-5-yl]-carbamic acid propyl ester.*

LC/MS (m/z) 264.9 ( $[M]^+$ ); RT = 2.36 (UV, ELSD) 93%, 99%.  $^1\text{H}$  NMR (DMSO- $d_6$ ): 0.89 (t, 3H), 1.59 (m, 2H), 2.99 (t, 2H), 3.50 (t, 2H), 3.97 (t, 2H), 5.96 (br. s, NH), 6.92 (s, 1H), 7.24 (s, 1H), 9.22 (s, 1H, NHCO).

5 *3,3-Dimethyl-N-(6-bromo-2,3-dihydro-1H-indol-5-yl)-butyramide.*

To a stirred solution of 3,3-dimethyl-N-[1-(2,2,2-trifluoroacetyl)-2,3-dihydro-1H-indol-5-yl]-butyramide (0.624 g, 1.90 mmol) in acetic acid (20 ml) bromine (0.195 ml, 1 eq.) was added. After 45 min more bromine (0.195 ml) was added. The reaction mixture was poured into solution of  $\text{Na}_2\text{SO}_3$  (5 g) in water (100 ml). The product N-[6-bromo-1-(2,2,2-trifluoroacetyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide  
10 was separated by filtration, washed with saturated aqueous  $\text{NaHCO}_3$  and water to furnish 0.555 g of colorless solid. Yield 71%. LC/MS (m/z) 409.0 ( $[M+1]^+$ ); RT = 3.38 (UV, ELSD) 97.5%, 85.5%.  $^1\text{H}$  NMR (DMSO- $d_6$ ): 1.05 (s, 9H), 2.25 (s, 2H), 3.22 (t, 2H), 4.30 (t, 2H), 7.55 (s, 1H), 8.25 (s, 1H), 9.37 (s, 1H, NHCO).

15 The solid (100 mg) was redissolved in methanol (10 ml) followed by addition of  $\text{K}_2\text{CO}_3$  (0.52 g) in water (5 ml). After stirring at 50°C for 5 min, the reaction mixture was poured into ice/water mixture and the title compound was isolated by filtration to give 0.057 g of colorless solid. Yield 75%. LC/MS (m/z) 313.0 ( $[M+1]^+$ ); RT = 1.71, (UV, ELSD) 97.5%, 98.9%.

20

**Preparation of intermediates of the general formula XVIII**

*N-(4-Chloro-2,3-dihydro-1H-indol-5-yl)-3,3-dimethylbutyramide.*

To a solution of crude N-[4-Chloro-1-(2,2,2-trifluoroacetyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide (ca. 100 mg) in MeOH (10 ml) a solution of  $\text{K}_2\text{CO}_3$  (0.5  
25 g) in water (2 ml) was added. The obtained mixture was heated at 50°C for 5 min and quenched with ethyl acetate and water. The organic solution was filtered via  $\text{SiO}_2$  (5 g) and evaporated in vacuo to furnish 20 mg of the title compound. The crude product was used in the next step without purification. LC/MS (m/z) 267.1 ( $[M+1]^+$ ); RT =  
30 1.61 (UV, ELSD) 45%, 78%.

*N*-(2,3-Dihydro-1H-indol-5-yl)-2-(4-fluoro-phenyl)-acetamide.

To a solution of 2-(4-Fluorophenyl)-N-[1-(2,2,2-trifluoroacetyl)-2,3-dihydro-1H-indol-5-yl]-acetamide (1.3 g, 3.55 mmol) in methanol (50 ml) K<sub>2</sub>CO<sub>3</sub> (7.6 g) in water (20 ml) was added. The reaction mixture was kept at 50°C for 5 min, poured into  
5 water and the title compound was separated by filtration. Yield 0.742 g, 77.4%. LC/MS (m/z) 271.0 ([M+1]<sup>+</sup>); RT = 1.42, (UV, ELSD) 94.5%, 98.7%. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 2.85 (t, 2H), 3.36 (t, 2H), 3.55 (s, 2H); 5.28 (br, 1H, NH); 6.41 (d, 1H); 7.06 (dd, 1H); 7.12 (t, 2H); 7.28 (d, 1H); 7.35 (dd, 2H); 9.75 (s, 1H, NHCO).

10 The following compound was prepared analogously from 3,3-dimethyl-N-[1-(2,2,2-trifluoroacetyl)-2,3-dihydro-1H-indol-5-yl]-butyramide:

*N*-(2,3-Dihydro-1H-indol-5-yl)-3,3-dimethyl-butyramide.

LC/MS (m/z) 232.9 ([M+1]<sup>+</sup>); RT = 1.43, (UV, ELSD) 94.4%, 88.1%. <sup>1</sup>H NMR  
15 (DMSO-d<sub>6</sub>): 1.00 (s, 9H); 2.09 (s, 2H); 2.84 (t, 2H); 3.37 (t, 2H); 5.25 (br, 1H, NH); 6.41 (d, 1H); 7.02 (dd, 1H); 7.29 (d, 1H); 9.34 (s, 1H, NHCO).

**Compounds of the invention**

20

**Example 1**

**1a** *N*-[4-Chloro-1-(4-trifluoromethylbenzyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide.

25 To a solution of N-(4-chloro-2,3-dihydro-1H-indol-5-yl)-3,3-dimethylbutyramide (10 mg), 4-trifluoromethylbenzaldehyde (0.06 ml) and acetic acid (0.03 ml) in methanol (0.3 ml) NaBH<sub>3</sub>CN (100 mg) was added. After 60 min the reaction mixture was partitioned between ethyl acetate and saturated aqueous NaHCO<sub>3</sub> solution. The organic layer was filtered via plug of SiO<sub>2</sub> (2 g), evaporated and purified by  
30 preparative LC/MS to give 11 mg of the title compound as colourless solid. LC/MS (m/z) 425.2 ([M+1]<sup>+</sup>); RT = 4.01, (UV, ELSD) 95%, 99%. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 1.03 (s, 9H), 2.16 (s, 2H), 2.97 (t, 2H), 3.40 (t, 2H), 4.41 (s, 2H), 6.48 (d, 1H), 7.03 (d, 1H), 7.56 (d, 2H), 7.72 (d, 2H), 9.12 (s, 1H, NHCO).

The following compound was prepared analogously using 5-chloro-2-thiophenecarboxaldehyde:

5 **1b** *N*-[4-Chloro-1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide.

LC/MS (m/z) 397.0 ( $[M+1]^+$ ); RT = 3.91, (UV, ELSD) 97%, 99%.  $^1\text{H}$  NMR (DMSO- $d_6$ ): 1.03 (s, 9H), 2.17 (s, 2H), 2.92 (t, 2H), 3.37 (t, 2H), 4.46 (s, 2H), 6.60 (d, 1H), 6.95 (d, 1H), 6.99 (d, 1H), 7.07 (d, 1H), 9.12 (s, 1H, NHCO).

10 **Example 2**

**2a** [1-(4-Fluorobenzyl)-2,3-dihydro-1H-indol-5-yl]-carbamic acid propyl ester.

To a cold (ice/water bath) solution of 1-(4-fluorobenzyl)-5-aminoindoline in acetonitrile (0.2 M, 0.15 ml) propyl chloroformate (0.02 ml or ca. 20 mg) was added  
15 followed by pyridine (0.03 ml). The reaction mixture was allowed to stand at room temperature for 60 min and evaporated in vacuo. The title compound was separated by preparative LC/MS, yield 5.8 mg, 59%. LC/MS (m/z) 329.1 ( $[M+1]^+$ ); RT = 2.68, (UV, ELSD) 94%, 99%.

20 The following compounds were obtained analogously from corresponding 5-aminoindolines and commercially available appropriate chloroformates, carbamyl chlorides, sulphonyl chlorides, acid chlorides, di-tert-butyl dicarbonate ( $\text{Boc}_2\text{O}$ ) or isocyanates, which are listed in the Table 1 below. Pyridine was used as a base in case of chloroformates, carbamyl chlorides, and sulphonyl chlorides. Triethylamine was  
25 used as a base in case of acid chlorides. No base was used in case of isocyanates and  $\text{Boc}_2\text{O}$ :

**2b** *N*-[1-(4-Fluorobenzyl)-2,3-dihydro-1H-indol-5-yl]-*C*-phenyl-methanesulfonamide.

LC/MS (m/z) 395.3 ( $[M-1]^+$ ); RT = 3.17, (UV, ELSD) 80%, 100%.

30

**2c** 4-Fluoro-*N*-[1-(4-fluorobenzyl)-2,3-dihydro-1H-indol-5-yl]-benzamide.

LC/MS (m/z) 365.4 ( $[M+1]^+$ ); RT = 2.90, (UV, ELSD) 96%, 100%.

**2d** *N*-[1-(4-Fluorobenzyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide.

LC/MS (m/z) 341.1 ( $[M+1]^+$ ); RT = 2.79, (UV, ELSD) 94%, 100%.

**2e** *N*-[1-(4-Fluorobenzyl)-2,3-dihydro-1H-indol-5-yl]-2-thiophen-2-ylacetamide.

5 LC/MS (m/z) 367.1 ( $[M+1]^+$ ); RT = 2.72, (UV, ELSD) 93%, 100%.

**2f** *N*-[1-(4-Fluorobenzyl)-2,3-dihydro-1H-indol-5-yl]-2-(4-fluorophenyl)-acetamide.

LC/MS (m/z) 379.3 ( $[M+1]^+$ ); RT = 2.82, (UV, ELSD) 95%, 100%.

10 **2g** 3-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-1,1-diisopropylurea.

LC/MS (m/z) 392.3 ( $[M+1]^+$ ); RT = 3.14, (UV, ELSD) 75%, 89%.

**2h** Morpholine-4-carboxylic acid [1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-amide.

15 LC/MS (m/z) 378.2 ( $[M+1]^+$ ); RT = 2.33, (UV, ELSD) 97%, 100%.

**2i** Pyrrolidine-1-carboxylic acid [1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-amide.

20 LC/MS (m/z) 362.0 ( $[M+1]^+$ ); RT = 2.48, (UV, ELSD) 83%, 99%.

**2j** [1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-carbamic acid 2-benzyloxyethyl ester.

LC/MS (m/z) 442.1 ( $[M]^+$ ); RT = 3.52, (UV, ELSD) 62%, 86%.

25

**2k** 3-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-1-methyl-1-propylurea.

LC/MS (m/z) 364.3 ( $[M+1]^+$ ); RT = 2.73, (UV, ELSD) 94%, 100%.

30 **2l** [1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-carbamic acid tert-butyl ester.

LC/MS (m/z) 364.3 ( $[M]^+$ ); RT = 3.50, (UV, ELSD) 97%, 100%.

**2m** *N*-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-C-phenyl-methanesulfonamide.

LC/MS (m/z) 418.2 ( $[M]^+$ ); RT = 3.44, (UV, ELSD) 98%, 100%.

5 **2n** *Butane-1-sulfonic acid* [1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-amide.

LC/MS (m/z) 384.1 ( $[M]^+$ ); RT = 3.43, (UV, ELSD) 98%, 100%.

10 **2o** *N*-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-4-fluorobenzamide.

LC/MS (m/z) 386.0 ( $[M]^+$ ); RT = 3.35, (UV, ELSD) 91%, 100%.

**2p** *N*-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-2,2-dimethylpropionamide.

15 LC/MS (m/z) 349.0 ( $[M+1]^+$ ); RT = 3.21, (UV, ELSD) 94%, 100%.

**2q** *N*-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-2-phenoxyacetamide.

LC/MS (m/z) 398.0 ( $[M]^+$ ); RT = 3.46, (UV, ELSD) 80%, 100%.

20

**2r** *N*-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide.

LC/MS (m/z) 362.1 ( $[M]^+$ ); RT = 3.34, (UV, ELSD) 84%, 99%.

25 **2s** *N*-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-butyramide.

LC/MS (m/z) 335.0 ( $[M+1]^+$ ); RT = 2.95, (UV, ELSD) 78%, 99%.

**2t** *Cyclopentanecarboxylic acid* [1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-amide.

30 LC/MS (m/z) 361.1 ( $[M+1]^+$ ); RT = 3.22, (UV, ELSD) 84%, 99%.

**2u** *N*-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-2-thiophen-2-ylacetamide.

LC/MS (m/z) 388.1 ( $[M]^+$ ); RT = 3.22, (UV, ELSD) 76%, 98%.

**2v** *N*-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-isonicotinamide.

LC/MS (m/z) 370.0 ( $[M+1]^+$ ); RT = 2.22, (UV, ELSD) 96%, 100%.

5

**2w** *N*-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-4-dimethylaminobenzamide.

LC/MS (m/z) 412.0 ( $[M+1]^+$ ); RT = 3.09, (UV, ELSD) 87%, 100%.

10 **2x** *N*-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-2-(4-fluorophenyl)-acetamide.

LC/MS (m/z) 401.0 ( $[M+1]^+$ ); RT = 3.31, (UV, ELSD) 84%, 100%.

15 **2y** *N*-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-6-trifluoromethylnicotinamide.

LC/MS (m/z) 437.1 ( $[M]^+$ ); RT = 3.46, (UV, ELSD) 90%, 99%.

**2z** 1-tert-Butyl-3-[1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-urea.

LC/MS (m/z) 364.3 ( $[M+1]^+$ ); RT = 2.80, (UV, ELSD) 97%, 100%.

20

**2aa** 1-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-3-ethylurea.

LC/MS (m/z) 335.1 ( $[M]^+$ ); RT = 2.34, (UV, ELSD) 96%, 100%.

**2ab** 1-Benzyl-3-[1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-urea.

25 LC/MS (m/z) 398.2 ( $[M+1]^+$ ); RT = 2.85, (UV, ELSD) 84%, 100%.

**2ac** 1-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-3-phenethylurea.

LC/MS (m/z) 411.9 ( $[M+1]^+$ ); RT = 3.00, (UV, ELSD) 87%, 97%.

30 **2ad** 1-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-3-thiophen-2-ylurea.

LC/MS (m/z) 390.0 ( $[M+1]^+$ ); RT = 3.01, (UV, ELSD) 94%, 92%.



**2ae** 1-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-3-thiophen-3-ylurea.

LC/MS (m/z) 390.2 ( $[M+1]^+$ ); RT = 2.98, (UV, ELSD) 96%, 100%.

5    **2af** [1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-carbamic acid propyl ester.

LC/MS (m/z) 351.2 ( $[M]^+$ ); RT = 3.48, (UV, ELSD) 94.0%, 98.0%.  $^1\text{H}$  NMR (DMSO- $d_6$ ): 0.92 (t, 3H); 1.62 (sextet, 2H); 2.84 (t, 2H); 3.22 (t, 2H), 3.89 (t, 2H), 4.38 (s, 2H), 6.59 (d, 1H); 6.92 (d, 1H); 6.98 (d, 1H); 7.07 (br. d (dd), 1H); 7.18 (br. s,

10    1H); 9.20 (br s, 1H, NH).

**Table 1.** Chloroformates, sulfonyl chlorides, carbamyl chlorides, acid chlorides, and isocyanates used in the preparation of compounds of the invention **2a – 2af**

Compound of the invention	Reagent	MW	Supplier	Catalog number
<b>2a</b>	n-PROPYL CHLOROFORMATE	122.55	Aldrich	24,946-7
<b>2b</b>	ALPHA-TOLUENESULFONYL CHLORIDE	190.649	Aldrich	15,971-9
<b>2c</b>	4-FLUOROBENZOYL CHLORIDE	158.559	Aldrich	11,994-6
<b>2d</b>	TERT-BUTYLACETYL CHLORIDE	134.605	Aldrich	B8,880-2
<b>2e</b>	THIOPHENE-2-ACETYL CHLORIDE	160.624	Aldrich	19,599-5
<b>2f</b>	4-FLUOROPHENYLACETYL CHLORIDE	172.585	Aldrich	46,695-6
<b>2g</b>	DIISOPROPYLCARBAMYL CHLORIDE	163.647	Aldrich	S31,027-1
<b>2h</b>	MORPHOLINE-4-CARBONYL CHLORIDE	149.576	Aldrich	34,829-5
<b>2i</b>	1-PYRROLIDINECARBONYL CHLORIDE	133.577	Aldrich	20,635-0
<b>2j</b>	2-BENZYLOXYETHYL CHLOROFORMATE	214.647	Aldrich	52,514-6
<b>2k</b>	N-ISOPROPYL-N-METHYL-CARBAMYL CHLORIDE	135.59	Lundbeck	C0005221
<b>2l</b>	DI-TERT-BUTYL DICARBONATE	218.247	Fluka	34660
<b>2m</b>	ALPHA-TOLUENESULFONYL CHLORIDE	190.649	Aldrich	15,971-9
<b>2n</b>	1-BUTANESULFONYL CHLORIDE	156.632	Aldrich	26,360-5
<b>2o</b>	4-FLUOROBENZOYL CHLORIDE	158.559	Aldrich	11,994-6
<b>2p</b>	PIVALOYL CHLORIDE	120.578	Aldrich	T7,260-5
<b>2q</b>	PHENOXYACETYL CHLORIDE	170.594	Aldrich	15,862-3
<b>2r</b>	TERT-BUTYLACETYL CHLORIDE	134.605	Aldrich	B8,880-2
<b>2s</b>	BUTYRYL CHLORIDE	106.551	Aldrich	10,961-4
<b>2t</b>	CYCLOPENTANECARBONYL CHLORIDE	132.589	Aldrich	32,831-6
<b>2u</b>	THIOPHENE-2-ACETYL CHLORIDE	160.624	Aldrich	19,599-5
<b>2v</b>	ISONICOTINOYL CHLORIDE HYDROCHLORIDE	178.018	Aldrich	22,875-3
<b>2w</b>	4-DIMETHYLAMINO BENZOYL CHLORIDE	183.637	Aldrich	52,611-8
<b>2x</b>	4-FLUOROPHENYLACETYL CHLORIDE	172.585	Aldrich	46,695-6
<b>2y</b>	6-(TRIFLUOROMETHYL)NICOTINOYL CHLORIDE	209.554	Fluoroche m	9368
<b>2z</b>	TERT-BUTYL ISOCYANATE	99.132	Aldrich	14,445-2
<b>2aa</b>	ETHYL ISOCYANATE	71.08	Aldrich	E3,330-0
<b>2ab</b>	BENZYL ISOCYANATE	133.149	Aldrich	22,726-9
<b>2ac</b>	PHENETHYL ISOCYANATE	147.176	Aldrich	45,617-9
<b>2ad</b>	2-THIENYL ISOCYANATE	125.151	Maybridge	CC 13006
<b>2ae</b>	3-THIENYL ISOCYANATE	125.151	Maybridge	CC 13106
<b>2af</b>	n-PROPYL CHLOROFORMATE	122.55	Aldrich	24,946-7

**Example 3**

**3a** *2,2-Dimethyl-N-[6-nitro-1-(4-trifluoromethylbenzyl)-2,3-dihydro-1H-indol-5-yl]-propionamide.*

- 5 To a stirred solution of 2,2-Dimethyl-N-(6-nitro-2,3-dihydro-1H-indol-5-yl)-propionamide (0.379 g, 1.44 mmol) in methanol (25 ml) 4-trifluoromethylbenzaldehyde (0.8 ml), acetic acid (0.8 ml) and a solution of NaBH<sub>3</sub>CN (0.8 g) in methanol (10 ml) were added in 4 portions during 3 hours until reaction completion. The obtained reaction mixture was concentrated in vacuo to a
- 10 small volume, quenched with saturated aqueous NaHCO<sub>3</sub> solution, and sonicated for several minutes. The title compound was separated by filtration to give 0.574g of red solid, yield 95%. LC/MS (m/z) 422.1 ([M+1]<sup>+</sup>); RT = 4.11, (UV, ELSD) 96%, 99%. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 1.20 (s, 9H), 3.04 (t, 2H), 3.42 (t, 2H), 4.49 (s, 2H), 7.09 (s, 1H), 7.48 (s, 1H), 7.57 (d, 2H), 7.73 (d, 2H), 9.62 (s, 1H, NHCO).

15

The following compounds were prepared analogously:

**3b** *N-[1-(5-Chlorothiophen-2-ylmethyl)-6-nitro-2,3-dihydro-1H-indol-5-yl]-2,2-dimethylpropionamide.*

- 20 LC/MS (m/z) 394.0 ([M+1]<sup>+</sup>); RT = 4.07, (UV, ELSD) 97%, 98%. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 1.20 (s, 9H), 3.00 (t, 2H), 3.41 (t, 2H), 4.55 (s, 2H), 6.97 (d, 1H), 7.01 (d, 1H), 7.22 (s, 1H), 7.47 (s, 1H), 9.62 (s, 1H, NHCO).

**3c** *2-(4-Fluorophenyl)-N-[6-nitro-1-(4-trifluoromethylbenzyl)-2,3-dihydro-1H-indol-5-yl]-acetamide.*

25

LC/MS (m/z) 474.2 ([M+1]<sup>+</sup>); RT = 3.89, (UV, ELSD) 80%, 97%.

**3d** *N-[1-(5-Chlorothiophen-2-ylmethyl)-6-nitro-2,3-dihydro-1H-indol-5-yl]-2-(4-fluorophenyl)-acetamide.*

- 30 LC/MS (m/z) 445.1 ([M]<sup>+</sup>); RT = 3.88, (UV, ELSD) 79.8%, 98.9%. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 2.98 (t, 2H); 3.41 (t, 2H), 3.63 (s, 2H), 4.54 (s, 2H), 6.96 (d, 1H); 7.0 (d, 1H); 7.16 (t, 2H); 7.17 (s, 1H); 7.31 (s, 1H); 7.34 (dd, 2H); 10.03 (s, 1H, NH).

**3e** *N*-[1-(5-Chlorothiophen-2-ylmethyl)-6-nitro-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide.

LC/MS (m/z) 407.1 ( $[M]^+$ ); RT = 4.07, (UV, ELSD) 72.4%, 98.7%.  $^1\text{H}$  NMR (DMSO- $d_6$ ): 1.01 (s, 9H); 2.16 (s, 2H); 2.99 (t, 2H); 3.41 (t, 2H), 4.54 (s, 2H), 6.98 (d, 1H); 7.01 (d, 1H); 7.14 (s, 1H); 7.23 (s, 1H); 9.76 (s, 1H, NH).

#### Example 4

**4a** *N*-[6-Amino-1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide.

To a solution of 3,3-dimethyl-N-(6-nitro-2,3-dihydro-1H-indol-5-yl)-butyramide (15 mg), 5-chloro-2-thiophenecarboxaldehyde (50 mg), and acetic acid (0.1 ml) in methanol (5 ml)  $\text{NaBH}_3\text{CN}$  (200 mg) was added. After 30 min the reaction mixture was concentrated in vacuo to a small volume and partitioned between ethyl acetate and water. The organic solution was washed with aqueous HCl (1 M) and saturated aqueous  $\text{NaHCO}_3$  and evaporated in vacuo.

The obtained residue was dissolved in tetrahydrofuran (10 ml) and acetic acid (2 ml) followed by addition of Zn powder (1 g). The obtained suspension was sonicated for 5 min, more Zn powder was added (0.5 g) and sonication continued for 2 min. The obtained suspension was filtered via a plug of  $\text{SiO}_2$  (2 g), evaporated, and the title compound was separated by preparative LC/MS to give 6.5 mg of colourless solid, yield 32%. LC/MS (m/z) 378.0 ( $[M+1]^+$ ); RT = 2.36, (UV, ELSD) 93%, 98%.  $^1\text{H}$  NMR (DMSO- $d_6$ ): 1.03 (s, 9H), 2.17 (s, 2H), 2.82 (t, 2H), 3.30 (t, 2H), 3.55 (very br. s,  $\text{NH}_2$  and  $\text{H}_2\text{O}$ ), 4.38 (s, 2H), 6.27 (s, 1H), 6.81 (s, 1H), 6.93 (d, 1H), 7.00 (d, 1H), 9.27 (br. s, 1H,  $\text{NHCO}$ ).

**4b** *N*-[6-Amino-1-(4-trifluoromethylbenzyl)-2,3-dihydro-1H-indol-5-yl]-2,2-dimethylpropionamide.

The title compound was prepared from the above 2,2-dimethyl-N-[6-nitro-1-(4-trifluoromethylbenzyl)-2,3-dihydro-1H-indol-5-yl]-propionamide (see **example 3**) by reduction with Zn powder as described above for 1-(5-Chlorothiophen-2-ylmethyl)-5-aminoindoline (see preparation of intermediates of the general formula **XXI**). The crude solid residue after filtration via  $\text{SiO}_2$  was treated with ethyl acetate and heptane

and the title compound was separated by filtration. Yield 0.375 g, 71%, colourless solid. LC/MS (m/z) 392.3 ( $[M+1]^+$ ); RT = 2.38, (UV, ELSD) 97%, 99%.  $^1\text{H}$  NMR (DMSO- $d_6$ ): 1.20 (s, 9H), 2.77 (t, 2H), 3.25 (t, 2H), 4.28 (s, 2H), 4.34 (s, 2H,  $\text{NH}_2$ ), 5.96 (s, 1H), 6.64 (s, 1H), 7.56 (d, 2H), 7.71 (d, 2H), 8.47 (s, 1H,  $\text{NHCO}$ ).

5

The following compounds were prepared analogously from corresponding 6-nitroindolines of the general formula XXV in two steps via reductive alkylation with appropriate aldehyde as described in the **example 3** followed by reduction with Zn powder as described above.

10

**4c** *N*-[6-Amino-1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-2,2-dimethylpropionamide.

LC/MS (m/z) 364.2 ( $[M+1]^+$ ); RT = 2.19, (UV, ELSD) 98%, 99%.  $^1\text{H}$  NMR (DMSO- $d_6$ ): 1.20 (s, 9H), 2.72 (t, 2H), 3.21 (t, 2H), 4.33 (s, 2H), 4.39 (br. s, 2H,  $\text{NH}_2$ ), 6.07 (s, 1H), 6.64 (s, 1H), 6.91 (d, 1H), 6.98 (d, 1H), 8.48 (s, 1H,  $\text{NHCO}$ ).

15

**4d** *N*-[6-Amino-1-(4-trifluoromethylbenzyl)-2,3-dihydro-1H-indol-5-yl]-2-(4-fluorophenyl)-acetamide.

LC/MS (m/z) 444.0 ( $[M+1]^+$ ); RT = 2.65, (UV, ELSD) 95%, 99%.  $^1\text{H}$  NMR (DMSO- $d_6$ ): 2.76 (t, 2H), 3.24 (t, 2H), 3.57 (s, 2H), 4.26 (s, 2H), 4.51 (br. s, 2H,  $\text{NH}_2$ ), 5.92 (s, 1H), 6.73 (s, 1H), 7.14 (t, 2H), 7.36 (dd, 2H), 7.55 (d, 2H), 7.71 (dd, 2H), 9.08 (s, 1H,  $\text{NHCO}$ ).

20

**4e** *N*-[6-Amino-1-(4-trifluoromethylbenzyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide.

LC/MS (m/z) 406.0 ( $[M+1]^+$ ); RT = 2.58, (UV, ELSD) 97.4%, 99.0%.  $^1\text{H}$  NMR (DMSO- $d_6$ ): 1.02 (s, 9H); 2.12 (s, 2H); 2.78 (t, 2H); 3.24 (t, 2H); 4.28 (s, 2H), 4.48 (s, 2H,  $\text{NH}_2$ ); 5.93 (s, 1H); 6.74 (s, 1H); 7.56 (d, 2H); 7.71 (d, 2H); 8.81 (s, 1H,  $\text{NH}$ ).

25

**4f** *N*-[6-Amino-1-(4-fluorobenzyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide.

30

LC/MS (m/z) 356.0 ( $[M+1]^+$ ); RT = 2.26, (UV, ELSD) 96.7%, 98.9%.  $^1\text{H}$  NMR (DMSO- $d_6$ ): 1.02 (s, 9H); 2.13 (s, 2H); 2.73 (t, 2H); 3.18 (t, 2H); 4.16 (s, 2H), 4.48 (s, 2H,  $\text{NH}_2$ ); 5.98 (s, 1H); 6.71 (s, 1H); 7.17 (t, 2H); 7.36 (dd, 2H); 8.80 (s, 1H, NH).

- 5    **4g** *N*-[6-Amino-1-(3-fluoro-4-trifluoromethylbenzyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide.

LC/MS (m/z) 424.0 ( $[M+1]^+$ ); RT = 2.58, (UV, ELSD) 92.0%, 98.8%.  $^1\text{H}$  NMR (DMSO- $d_6$ ): 1.01 (s, 9H); 2.12 (s, 2H); 2.79 (t, 2H); 3.28 (t, 2H); 4.27 (s, 2H), 4.48 (s, 2H,  $\text{NH}_2$ ); 5.90 (s, 1H); 6.75 (s, 1H); 7.39 (d, 1H); 7.45 (d, 1H); 7.78 (t, 1H); 8.80 (s, 1H, NH).

### Example 5

**5a** *N*-[1-(5-Chlorothiophen-2-ylmethyl)-1H-indol-5-yl]-3,3-dimethylbutyramide.

- 15    To a solution of *N*-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide (20 mg) in dimethylsulfoxide- $d_6$  (0.6 ml) 2,3,5,6-tetrachloro-[1,4]benzoquinone (65 mg) was added. The obtained mixture was heated at 70°C for 5 min, allowed to cool and poured into aqueous  $\text{NaHSO}_3$  solution (1 g in 5 ml) followed by addition of 25% aqueous  $\text{NH}_3$  (5 ml) and 10% aqueous  $\text{NaOH}$  (5 ml).
- 20    The mixture was extracted with  $\text{CH}_2\text{Cl}_2$  (3 x 10 ml), the combined organic solution was washed with water and 1M  $\text{HCl}$ , filtered via plug of  $\text{SiO}_2$  (10 g) and eluted with ethyl acetate/heptane (1:1). The crude product after evaporation was purified by preparative LC/MS to give 5 mg of the title compound as colourless solid. LC/MS (m/z) 361.1 ( $[M+1]^+$ ); RT = 3.43, (UV, ELSD) 96%, 99%.  $^1\text{H}$  NMR (DMSO- $d_6$ ): 1.03
- 25    (s, 9H), 2.16 (s, 2H), 5.51 (s, 2H), 6.41 (d, 1H), 6.95 (d, 1H), 6.98 (d, 1H), 7.21 (dd, 1H), 7.41 (d, 1H), 7.44 (d, 1H), 7.87 (d, 1H), 9.60 (s, 1H,  $\text{NHCO}$ ).

### Example 6

- 30    **6a**        *N*-[6-Bromo-1-(4-trifluoromethylbenzyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide.

The title compound was prepared from 3,3-dimethyl-*N*-(6-bromo-2,3-dihydro-1H-indol-5-yl)-butyramide (25 mg) and 4-trifluoromethylbenzaldehyde (64 mg) as

described in example 1. Yield 16.7 mg. LC/MS (m/z) 469.1 ( $[M+1]^+$ ); RT = 3.99, (UV, ELSD) 97.4%, 95.1%.

The following compound was prepared analogously using appropriate aldehyde:

5

**6b** *N*-[6-Bromo-1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide.

Yield 16.8 mg. LC/MS (m/z) 443.1 ( $[M+1]^+$ ); RT = 3.92, (UV, ELSD) 96.3%, 94.9%.

**Example 7**

10

General procedure: To a mixture of aldehyde (25 mg or 0.025 ml) and appropriate indoline (7 mg) in methanol (0.5 ml) NaBH<sub>3</sub>CN (20 mg) in methanol (0.2 ml) was added followed by acetic acid (0.05 ml). The obtained solution or suspension was sonicated for 5 min and then kept at 50°C for 60 min followed by evaporation in vacuo. The residue was dissolved in DMSO (2.5 ml) and water (100 ml). The product was isolated by preparative LC/MS.

15

The following compounds were prepared accordingly from either N-(2,3-Dihydro-1H-indol-5-yl)-3,3-dimethyl-butylamide (compounds **7a-7z**) or from N-(2,3-Dihydro-1H-indol-5-yl)-2-(4-fluoro-phenyl)-acetamide (compounds **7aa-7aw**) and the appropriate aldehyde (see Table 2 below):

20

**7a** *N*-[1-(4-Chlorobenzyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide.

LC/MS (m/z) 357.2 ( $[M+1]^+$ ); RT = 3.17, (UV, ELSD) 77.1%, 97.2%.

25

**7b** 3,3-Dimethyl-N-[1-(4-trifluoromethylbenzyl)-2,3-dihydro-1H-indol-5-yl]-butyramide.

LC/MS (m/z) 391.4 ( $[M+1]^+$ ); RT = 3.44, (UV, ELSD) 91.8%, 99.6%.

30

**7c** *N*-[1-(4-Isopropylbenzyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide.

LC/MS (m/z) 365.4 ( $[M+1]^+$ ); RT = 3.13, (UV, ELSD) 89.4%, 99.2%.

- 7d** *N*-[1-(3-Fluoro-4-trifluoromethylbenzyl)-2,3-dihydro-1*H*-indol-5-yl]-3,3-dimethylbutyramide.  
LC/MS (m/z) 409.2 ( $[M+1]^+$ ); RT = 3.60, (UV, ELSD) 69.6%, 98.2%.
- 5 **7e** *N*-[1-(6-Chlorobenzo[1,3]dioxol-5-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-3,3-dimethylbutyramide.  
LC/MS (m/z) 401.0 ( $[M+1]^+$ ); RT = 3.22, (UV, ELSD) 94.6%, 99.5%.
- 10 **7f** *N*-[1-(3,5-Dimethyl-1-phenyl-1*H*-pyrazol-4-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-3,3-dimethylbutyramide.  
LC/MS (m/z) 417.2 ( $[M+1]^+$ ); RT = 2.23, (UV, ELSD) 93.3%, 98.9%.
- 15 **7g** *N*-[1-(2-Chloro-5-trifluoromethylbenzyl)-2,3-dihydro-1*H*-indol-5-yl]-3,3-dimethylbutyramide.  
LC/MS (m/z) 425.2 ( $[M+1]^+$ ); RT = 3.79, (UV, ELSD) 75.4%, 98.4%.
- 7h** *N*-{1-[5-(4-Chlorophenoxy)-1,3-dimethyl-1*H*-pyrazol-4-ylmethyl]-2,3-dihydro-1*H*-indol-5-yl}-3,3-dimethylbutyramide.  
LC/MS (m/z) 467.3 ( $[M+1]^+$ ); RT = 2.71, (UV, ELSD) 95.2%, 99.8%.
- 20 **7i** 3,3-Dimethyl-*N*-[1-(6-*p*-tolylloxy-pyridin-3-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-butyramide.  
LC/MS (m/z) 430.2 ( $[M+1]^+$ ); RT = 3.15, (UV, ELSD) 76.6%, 97.9%.
- 25 **7k** *N*-{1-[6-(4-Chlorophenylsulfanyl)-pyridin-3-ylmethyl]-2,3-dihydro-1*H*-indol-5-yl}-3,3-dimethylbutyramide.  
LC/MS (m/z) 466.0 ( $[M+1]^+$ ); RT = 3.45, (UV, ELSD) 69.3%, 97.1%.
- 30 **7l** *N*-{1-[6-(4-Cyanophenoxy)-pyridin-3-ylmethyl]-2,3-dihydro-1*H*-indol-5-yl}-3,3-dimethylbutyramide.  
LC/MS (m/z) 441.4 ( $[M+1]^+$ ); RT = 2.98, (UV, ELSD) 79.8%, 98.9%.



**7m** 3,3-Dimethyl-N-[1-(6-trifluoromethylpyridin-3-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-butyramide.

LC/MS (m/z) 392.4 ( $[M+1]^+$ ); RT = 3.10, (UV, ELSD) 82.5%, 99.5%.

5 **7p** 3,3-Dimethyl-N-[1-(3-methyl-benzo[b]thiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-butyramide.

LC/MS (m/z) 393.3 ( $[M+1]^+$ ); RT = 3.56, (UV, ELSD) 72.8%, 97.9%.

10 **7q** N-[1-(6-Fluoro-4H-benzo[1,3]dioxin-8-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide.

LC/MS (m/z) 399.2 ( $[M+1]^+$ ); RT = 2.75, (UV, ELSD) 84.9%, 99.3%.

**7s** 3,3-Dimethyl-N-[1-(6-phenoxy-pyridin-3-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-butyramide.

15 LC/MS (m/z) 416.2 ( $[M+1]^+$ ); RT = 2.98, (UV, ELSD) 67.5%, 96.5%.

**7u** 3,3-Dimethyl-N-[1-(3-methyl-5-phenyl-isoxazol-4-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-butyramide.

LC/MS (m/z) 404.4 ( $[M+1]^+$ ); RT = 3.24, (UV, ELSD) 97.6%, 99.9%.

20

**7v** N-(1-Benzo[b]thiophen-2-ylmethyl-2,3-dihydro-1H-indol-5-yl)-3,3-dimethylbutyramide.

LC/MS (m/z) 379.3 ( $[M+1]^+$ ); RT = 3.44, (UV, ELSD) 71.8%, 97.6%.

25 **7w** N-{1-[1-(4-Fluorophenyl)-5-methyl-1H-pyrazol-4-ylmethyl]-2,3-dihydro-1H-indol-5-yl}-3,3-dimethylbutyramide.

LC/MS (m/z) 421.4 ( $[M+1]^+$ ); RT = 2.24, (UV, ELSD) 79.5%, 98.9%.

30 **7y** 3,3-Dimethyl-N-[1-(5-methylthiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-butyramide.

LC/MS (m/z) 343.1 ( $[M+1]^+$ ); RT = 2.84, (UV, ELSD) 59.9%, 89.1%.

**7z** 3,3-Dimethyl-N-[1-(4-pyrrol-1-yl-benzyl)-2,3-dihydro-1H-indol-5-yl]-butyramide.

LC/MS (m/z) 388.3 ( $[M+1]^+$ ); RT = 3.03, (UV, ELSD) 81.1%, 99.5%.

**7aa** *N*-[1-(4-Chlorobenzyl)-2,3-dihydro-1H-indol-5-yl]-2-(4-fluorophenyl)-acetamide.

LC/MS (m/z) 395.1 ( $[M+1]^+$ ); RT = 2.99, (UV, ELSD) 93.4%, 93.1%.

5

**7ab** 2-(4-Fluorophenyl)-*N*-[1-(4-trifluoromethylbenzyl)-2,3-dihydro-1H-indol-5-yl]-acetamide.

LC/MS (m/z) 429.1 ( $[M+1]^+$ ); RT = 3.24, (UV, ELSD) 76.6%, 87.4%.

10 **7ac** 2-(4-Fluorophenyl)-*N*-[1-(4-isopropylbenzyl)-2,3-dihydro-1H-indol-5-yl]-acetamide.

LC/MS (m/z) 403.1 ( $[M+1]^+$ ); RT = 2.96, (UV, ELSD) 91.5%, 83.7%.

15 **7ad** 2-(4-Fluorophenyl)-*N*-[1-(3-fluoro-4-trifluoromethylbenzyl)-2,3-dihydro-1H-indol-5-yl]-acetamide.

LC/MS (m/z) 447.2 ( $[M+1]^+$ ); RT = 3.36, (UV, ELSD) 79.3%, 90.4%.

**7ae** *N*-[1-(6-Chlorobenzo[1,3]dioxol-5-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-2-(4-fluorophenyl)-acetamide.

20 LC/MS (m/z) 439.0 ( $[M+1]^+$ ); RT = 3.02, (UV, ELSD) 94.0%, 92.2%.

**7af** *N*-[1-(3,5-Dimethyl-1-phenyl-1H-pyrazol-4-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-2-(4-fluorophenyl)-acetamide.

LC/MS (m/z) 455.0 ( $[M+1]^+$ ); RT = 2.14, (UV, ELSD) 97.1%, 91.4%.

25

**7ag** *N*-[1-(2-Chloro-5-trifluoromethylbenzyl)-2,3-dihydro-1H-indol-5-yl]-2-(4-fluorophenyl)-acetamide.

LC/MS (m/z) 463.1 ( $[M+1]^+$ ); RT = 3.53, (UV, ELSD) 98.3%, 95.3%.

30 **7ah** *N*-{1-[5-(4-Chlorophenoxy)-1,3-dimethyl-1H-pyrazol-4-ylmethyl]-2,3-dihydro-1H-indol-5-yl}-2-(4-fluorophenyl)-acetamide.

LC/MS (m/z) 503.3 ( $[M-1]^+$ ); RT = 2.55, (UV, ELSD) 92.3%, 90.5%.

**7ai** *N*-{1-[6-(4-Cyanophenoxy)-pyridin-3-ylmethyl]-2,3-dihydro-1*H*-indol-5-yl}-2-(4-fluorophenyl)-acetamide.

LC/MS (*m/z*) 479.1 ( $[M+1]^+$ ); RT = 2.81, (UV, ELSD) 84.1%, 87.1%.

5 **7al** 2-(4-Fluorophenyl)-*N*-[1-(3-methyl-benzo[*b*]thiophen-2-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-acetamide.

LC/MS (*m/z*) 431.2 ( $[M+1]^+$ ); RT = 3.34, (UV, ELSD) 60.2%, 88.8%.

10 **7am** *N*-[1-(6-Fluoro-4*H*-benzo[1,3]dioxin-8-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-2-(4-fluorophenyl)-acetamide.

LC/MS (*m/z*) 437.0 ( $[M+1]^+$ ); RT = 2.62, (UV, ELSD) 79.7%, 83.6%.

**7ao** 2-(4-Fluorophenyl)-*N*-[1-(6-phenoxy-pyridin-3-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-acetamide.

15 LC/MS (*m/z*) 454.3 ( $[M+1]^+$ ); RT = 2.81, (UV, ELSD) 77.9%, 83.1%.

**7ar** *N*-(1-Benzo[*b*]thiophen-2-ylmethyl-2,3-dihydro-1*H*-indol-5-yl)-2-(4-fluorophenyl)-acetamide.

LC/MS (*m/z*) 417.2 ( $[M+1]^+$ ); RT = 3.22, (UV, ELSD) 80.2%, 89.4%.

20

**7as** 2-(4-Fluorophenyl)-*N*-{1-[1-(4-fluorophenyl)-5-methyl-1*H*-pyrazol-4-ylmethyl]-2,3-dihydro-1*H*-indol-5-yl}-acetamide.

LC/MS (*m/z*) 459.3 ( $[M+1]^+$ ); RT = 2.15, (UV, ELSD) 91.5%, 88.7%.

25 **7au** 2-(4-Fluorophenyl)-*N*-[1-(5-methylthiophen-2-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-acetamide.

LC/MS (*m/z*) 381.1 ( $[M+1]^+$ ); RT = 2.71, (UV, ELSD) 76.6%, 82.9%.

**7av** 2-(4-Fluorophenyl)-*N*-[1-(4-pyrrol-1-yl-benzyl)-2,3-dihydro-1*H*-indol-5-yl]-acetamide.

30

LC/MS (*m/z*) 426.1 ( $[M+1]^+$ ); RT = 2.87, (UV, ELSD) 92.2%, 88.2%.

**Table 2.** Aldehydes used in the preparation of compounds of the invention 7a – 7av

Compound of the invention	Aldehyde	MW	Supplier	Catalog number
7a	4-CHLOROBENZALDEHYDE	140.568	Aldrich	
7b	4-(TRIFLUOROMETHYL)- BENZALDEHYDE	174.12	Aldrich	
7c	4-ISOPROPYLBENZALDEHYDE	148.204	Aldrich	
7d	3-FLUORO-4-(TRIFLUOROMETHYL)- BENZALDEHYDE	192.111	ABCR	AV20008
7e	6-CHLOROPIPERONAL	184.577	ABCR	AV13607
7f	3,5-DIMETHYL-1-PHENYLPYRAZOLE- 4-CARBOXALDEHYDE	200.24	Acros Organics	40852-0050
7g	2-CHLORO-5-(TRIFLUOROMETHYL)- BENZALDEHYDE	208.566	Aldrich	37,682-5
7h	5-(4-CHLOROPHENOXY)-1,3- DIMETHYL-1H-PYRAZOLE-4- CARBALDEHYDE	250.684	Bionet Research	11F-431S
7j	6-(4-METHYLPHENOXY)- NICOTINALDEHYDE	213.235	Bionet Research	5L-355S
7k	6-[(4-CHLOROPHENYL)SULFANYL]- NICOTINALDEHYDE	249.72	Bionet Research	5L-356S
7l	4-[(5-FORMYL-2-PYRIDINYLOXY)- BENZENECARBONITRILE	224.218	Bionet Research	6L-309S
7m	6-(TRIFLUOROMETHYL)PYRIDINE-3- CARBOXALDEHYDE	175.109	Fluoroche m	9397
7p	3-METHYLBENZO[B]THIOPHENE-2- CARBOXALDEHYDE	176.238	ABCR	AV11375
7q	5-FLUORO-4H-1,3-BENZODIOXINE-5- CARBALDEHYDE	182.149	Maybridge	CC 01904
7s	6-PHENOXYNICOTINALDEHYDE	199.208	Maybridge	CC 19604
7u	3-METHYL-5-PHENYL-4- ISOXAZOLECARBALDEHYDE	187.197	Maybridge	CC 20304
7v	1-BENZOTHIOPHENE-2- CARBALDEHYDE	162.211	Specs	942/2503463 9
7w	1-(4-FLUOROPHENYL)-5-METHYL-1H- PYRAZOLE-4-CARBALDEHYDE	204.203	Maybridge	MO 00310
7y	5-METHYL-2- THIOPHENECARBOXALDEHYDE	126.178	Aldrich	M8,441-0
7z	4-(1H-PYRROL-1-YL)BENZALDEHYDE	171.198	Maybridge	N/A
7aa	4-CHLOROBENZALDEHYDE	140.568	Aldrich	
7ab	4-(TRIFLUOROMETHYL)- BENZALDEHYDE	174.12	Aldrich	
7ac	4-ISOPROPYLBENZALDEHYDE	148.204	Aldrich	
7ad	3-FLUORO-4-(TRIFLUOROMETHYL)- BENZALDEHYDE	192.111	ABCR	AV20008

7ae	6-CHLOROPIPERONAL	184.577	ABCR	AV13607
7af	3,5-DIMETHYL-1-PHENYLPYRAZOLE-4-CARBOXALDEHYDE	200.24	Acros Organics	40852-0050
7ag	2-CHLORO-5-(TRIFLUOROMETHYL)BENZALDEHYDE	208.566	Aldrich	37,682-5
7ah	5-(4-CHLOROPHENOXY)-1,3-DIMETHYL-1H-PYRAZOLE-4-CARBALDEHYDE	250.684	Bionet Research	11F-431S
7ai	4-[(5-FORMYL-2-PYRIDINYL)OXY]BENZENECARBONITRILE	224.218	Bionet Research	6L-309S
7al	3-METHYLBENZO[B]THIOPHENE-2-CARBOXALDEHYDE	176.238	ABCR	AV11375
7am	5-FLUORO-4H-1,3-BENZODIOXINE-5-CARBALDEHYDE	182.149	Maybridge	CC 01904
7ao	6-PHENOXYNICOTINALDEHYDE	199.208	Maybridge	CC 19604
7ar	1-BENZOTHIOPHENE-2-CARBALDEHYDE	162.211	Specs	942/2503463 9
7as	1-(4-FLUOROPHENYL)-5-METHYL-1H-PYRAZOLE-4-CARBALDEHYDE	204.203	Maybridge	MO 00310
7au	5-METHYL-2-THIOPHENECARBOXALDEHYDE	126.178	Aldrich	M8,441-0
7av	4-(1H-PYRROL-1-YL)BENZALDEHYDE	171.198	Maybridge	N/A

**In vitro and in vivo testing**

The compounds of the invention have been tested and shown effect in one or more of the below models:

5

**Relative efflux through the KCNQ2 channel.**

This exemplifies a KCNQ2 screening protocol for evaluating compounds of the present invention. The assay measures the relative efflux through the KCNQ2 channel, and was carried out according to a method described by Tang et al. (Tang, 10 W. et. al., *J. Biomol. Screen.* **2001**, 6, 325-331) for hERG potassium channels with the modifications described below.

An adequate number of CHO cells stably expressing voltage-gated KCNQ2 channels were plated at a density sufficient to yield a mono-confluent layer on the day of the 15 experiment. Cells were seeded on the day before the experiment and loaded with 1  $\mu\text{Ci/ml}$  [ $^{86}\text{Rb}$ ] over night. On the day of the experiment cells were washed with a HBSS-containing buffer. Cells were pre-incubated with drug for 30 minutes and the  $^{86}\text{Rb}^+$  efflux was stimulated by a submaximal concentration of 15 mM KCl in the continued presence of drug for additional 30 minutes. After a suitable incubation 20 period, the supernatant was removed and counted in a liquid scintillation counter (Tricarb). Cells were lysed with 2 mM NaOH and the amount of  $^{86}\text{Rb}^+$  was counted. The relative efflux was calculated  $((\text{CPM}_{\text{super}}/(\text{CPM}_{\text{super}} + \text{CPM}_{\text{cell}}))_{\text{Cmpd}}/(\text{CPM}_{\text{super}}/(\text{CPM}_{\text{super}} + \text{CPM}_{\text{cell}}))_{15\text{mM KCl}}) * 100 - 100$ .

25 The compounds of the invention have an  $\text{EC}_{50}$  of less than 20000nM, in most cases less than 2000 nM and in many cases less than 200 nM. Accordingly, the compounds of the invention are considered to be useful in the treatment of diseases associated with the KCNQ family potassium channels.

**30 Electrophysiological patch-clamp recordings.**

Voltage-activated KCNQ2 currents were recorded from mammalian CHO cells by use of conventional patch-clamp recordings techniques in the whole-cell patch-clamp configuration (Hamill OP et.al. *Pflügers Arch* **1981**; 391: 85-100). CHO cells with

stable expression of voltage-activated KCNQ2 channels were grown under normal cell culture conditions in CO<sub>2</sub> incubators and used for electrophysiological recordings 1-7 days after plating. KCNQ2 potassium channels were activated by voltage steps up to + 80 mV in increments of 5-20 mV (or with a ramp protocol) from a membrane holding potential between - 100 mV and - 40 mV (Tatulian L et al. *J Neuroscience* 2001; 21 (15): 5535-5545). The electrophysiological effects induced by the compounds were evaluated on various parameters of the voltage-activated KCNQ2 current. Especially effects on the activation threshold for the current and on the maximum induced current were studied.

10

Some of the compounds of the invention have been tested in this test. A left-ward shift of the activation threshold or an increase in the maximum induced potassium current is expected to decrease the activity in neuronal networks and thus make the compounds useful in diseases with increased neuronal activity - like epilepsy.

15

#### **Maximum electroshock**

The test was conducted in groups of male mice using corneal electrodes and administering a square wave current of 26mA for 0.4 seconds in order to induce a convulsion characterised by a tonic hind limb extension (Wlaz et al. *Epilepsy Research* 1998, 30, 219-229).

20

#### **Pilocarpine induced seizures**

Pilocarpine induced seizures are induced by intraperitoneal injection of pilocarpine 250mg/kg to groups of male mice and observing for seizure activity resulting in loss of posture within a period of 30 minutes (Starr et al. *Pharmacology Biochemistry and Behavior* 1993, 45, 321-325).

25

#### **Electrical seizure -threshold test**

A modification of the up-and-down method (Kimball et al. *Radiation Research* 1957, 1-12) was used to determine the median threshold to induce tonic hind-limb extension in response to corneal electroshock in groups of male mice. The first mouse of each group received an electroshock at 14 mA, (0.4 s, 50 Hz) and was observed for seizure activity. If a seizure was observed the current was reduced by 1 mA for the next

30

mouse, however, if no seizure was observed then the current was increased by 1 mA. This procedure was repeated for all 15 mice in the treatment group.

#### **Chemical seizure -threshold test**

- 5 The threshold dose of pentylenetetrazole required to induce a clonic convulsion was measured by timed infusion of pentylenetetrazole (5mg / mL at 0.5 mL/minute) into a lateral tail vein of groups of male mice (Nutt et al. *J Pharmacy and Pharmacology* 1986, 38, 697-698).

#### **10 Amygdala kindling**

- Rats underwent surgery to implantation of tri-polar electrodes into the dorsolateral amygdala. After surgery the animals were allowed to recover before the groups of rats received either varying doses of test compound or the drug's vehicle. The animals were stimulated with their initial after discharge threshold + 25  $\mu$ A daily for 3-5 weeks and on each occasion seizure severity, seizure duration, and duration of electrical after discharge were noted. (Racine. *Electroencephalography and Clinical Neurophysiology* 1972, 32, 281-294).

#### **Side effects**

- 20 Central nervous system side-effects were measured by measuring the time mice would remain on rotarod apparatus (Capacio et al. *Drug and Chemical Toxicology* 1992, 15, 177-201); or by measuring their locomotor activity by counting the number of infra-red beams crossed in a test cage (Watson et al. *Neuropharmacology* 1997, 36, 1369-1375). Hypothermic actions on the animals core body temperature of the compound were measured by either rectal probe or implanted radiotelemetry transmitters capable of measuring temperature (Keeney et al. *Physiology and Behaviour* 2001, 74, 177-184).


#### **Pharmacokinetics**

- 30 The pharmacokinetic properties of the compounds were determined via. i.v. and p.o. dosing to Sprague Dawley rats, and, thereafter, drawing blood samples over 20 hours. Plasma concentrations were determined with LC/MS/MS.



## Claims

1 A substituted indoline or indole derivative of the general formula I



(I)

wherein

the dotted line represents an optional bond;

$R^1$  and  $R^{1'}$  are independently selected from the group consisting of hydrogen,  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, hydroxy- $C_{1-6}$ -alk(en/yn)yl, hydroxy- $C_{3-8}$ -cycloalk(en)yl, hydroxy- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, halo- $C_{1-6}$ -alk(en/yn)yl, halo- $C_{3-8}$ -cycloalk(en)yl, halo- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, cyano- $C_{1-6}$ -alk(en/yn)yl, cyano- $C_{3-8}$ -cycloalk(en)yl and cyano- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl; or

$R^1$  and  $R^{1'}$  together with the carbon atom to which they are attached form a 3-8 membered saturated or unsaturated ring which optionally contains 1 or 2 heteroatoms;

$s$  is 0 or 1;

$U$  is O,  $NR^{11}$ , S,  $SO_2$ ,  $SO_2NR^{11}$ , CO-O or CO- $NR^{11}$ ; wherein  $R^{11}$  is selected from the group consisting of hydrogen,  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl; or  $R^2$  and  $R^{11}$  together with the nitrogen atom to

which they are attached form a 4-8 membered saturated or unsaturated ring which optionally contains 1, 2 or 3 further heteroatoms;

**R**<sup>2</sup> is selected from the group consisting of hydrogen, C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, Ar, Ar-C<sub>1-6</sub>-alk(en/yn)yl, Ar-C<sub>3-8</sub>-cycloalk(en)yl, Ar-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, acyl, hydroxy-C<sub>1-6</sub>-alk(en/yn)yl, hydroxy-C<sub>3-8</sub>-cycloalk(en)yl, hydroxy-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, halogen, halo-C<sub>1-6</sub>-alk(en/yn)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, cyano, cyano-C<sub>1-6</sub>-alk(en/yn)yl, cyano-C<sub>3-8</sub>-cycloalk(en)yl, cyano-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, -NO<sub>2</sub>, NR<sup>10</sup>R<sup>10'</sup>-C<sub>1-6</sub>-alk(en/yn)yl, NR<sup>10</sup>R<sup>10'</sup>-C<sub>3-8</sub>-cycloalk(en)yl and NR<sup>10</sup>R<sup>10'</sup>-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl; wherein

**R**<sup>10</sup> and **R**<sup>10'</sup> are independently selected from the group consisting of hydrogen, C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, hydroxy-C<sub>1-6</sub>-alk(en/yn)yl, hydroxy-C<sub>3-8</sub>-cycloalk(en)yl, hydroxy-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, halo-C<sub>1-6</sub>-alk(en/yn)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, cyano-C<sub>1-6</sub>-alk(en/yn)yl, cyano-C<sub>3-8</sub>-cycloalk(en)yl and cyano-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, or

**R**<sup>10</sup> and **R**<sup>10'</sup> together with the nitrogen atom to which they are attached form a 4-8 membered saturated or unsaturated ring which optionally contains 1, 2 or 3 further heteroatoms;

with the proviso that when **R**<sup>2</sup> is NO<sub>2</sub>, halogen or cyano then s is 0; and

with the proviso that when **R**<sup>2</sup> is a hydrogen atom or acyl and s is 1 then **U** is NR<sup>11</sup>, O or S;

wherein the group -(U)<sub>s</sub>-**R**<sup>2</sup> is linked to position 4 or 6 of the indole or indoline;

**q** is 0 or 1;

**Z** is O or S;

**X** is CO or SO<sub>2</sub>; with the proviso that **q** is 0 when **X** is SO<sub>2</sub>;

$R^3$  is selected from the group consisting of  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl, heterocycloalk(en)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl,  $C_{1-6}$ -alk(en/yn)yl- $C_{3-8}$ -cycloalk(en)yl,  $C_{1-6}$ -alk(en/yn)yl-heterocycloalk(en)yl, Ar, Ar- $C_{1-6}$ -alk(en/yn)yl, Ar- $C_{3-8}$ -cycloalk(en)yl, Ar-heterocycloalk(en)yl, Ar- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, Ar- $C_{1-6}$ -alk(en/yn)yl- $C_{3-8}$ -cycloalk(en)yl, Ar- $C_{1-6}$ -alk(en/yn)yl-heterocycloalk(en)yl,  $C_{1-6}$ -alk(en/yn)yl-oxy- $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl-oxy- $C_{1-6}$ -alk(en/yn)yl,  $C_{1-6}$ -alk(en/yn)yl-oxy- $C_{3-8}$ -cycloalk(en)yl,  $C_{1-6}$ -alk(en/yn)yl-oxy-heterocycloalk(en)yl, Ar-oxy- $C_{1-6}$ -alk(en/yn)yl, Ar- $C_{1-6}$ -alk(en/yn)yl-oxy- $C_{1-6}$ -alk(en/yn)yl,  $C_{1-6}$ -alk(en/yn)yl-oxy-carbonyl- $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl-oxy-carbonyl- $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl-oxy-carbonyl- $C_{1-6}$ -alk(en/yn)yl, hydroxy- $C_{1-6}$ -alk(en/yn)yl, hydroxy- $C_{3-8}$ -cycloalk(en)yl, hydroxy-heterocycloalk(en)yl, hydroxy- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, hydroxy- $C_{1-6}$ -alk(en/yn)yl- $C_{3-8}$ -cycloalk(en)yl, hydroxy- $C_{1-6}$ -alk(en/yn)yl-heterocycloalk(en)yl, halo- $C_{1-6}$ -alk(en/yn)yl, halo- $C_{3-8}$ -cycloalk(en)yl, halo-heterocycloalk(en)yl, halo- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, halo- $C_{1-6}$ -alk(en/yn)yl- $C_{3-8}$ -cycloalk(en)yl, halo- $C_{1-6}$ -alk(en/yn)yl-heterocycloalk(en)yl, halo- $C_{1-6}$ -alk(en/yn)yl-Ar, halo- $C_{3-8}$ -cycloalk(en)yl-Ar, halo- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl-Ar, halo- $C_{1-6}$ -alk(en/yn)yl- $C_{3-8}$ -cycloalk(en)yl-Ar, cyano- $C_{1-6}$ -alk(en/yn)yl, cyano- $C_{3-8}$ -cycloalk(en)yl, cyano-heterocycloalk(en)yl, cyano- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, cyano- $C_{1-6}$ -alk(en/yn)yl- $C_{3-8}$ -cycloalk(en)yl, cyano- $C_{1-6}$ -alk(en/yn)yl-heterocycloalk(en)yl, acyl- $C_{1-6}$ -alk(en/yn)yl, acyl- $C_{3-8}$ -cycloalk(en)yl, acyl-heterocycloalk(en)yl, acyl- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, acyl- $C_{1-6}$ -alk(en/yn)yl- $C_{3-8}$ -cycloalk(en)yl, acyl- $C_{1-6}$ -alk(en/yn)yl-heterocycloalk(en)yl and  $-NR^{12}R^{12'}$ , optionally substituted  $NR^{12}R^{12'}$ - $C_{1-6}$ -alk(en/yn)yl, optionally substituted  $NR^{12}R^{12'}$ - $C_{3-8}$ -cycloalk(en)yl, optionally substituted  $NR^{12}R^{12'}$ - $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl; wherein  $R^{12}$  and  $R^{12'}$  are independently selected from the group consisting of hydrogen,  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, Ar, Ar- $C_{1-6}$ -alk(en/yn)yl, Ar- $C_{3-8}$ -cycloalk(en)yl, Ar- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, hydroxy- $C_{1-6}$ -alk(en/yn)yl, hydroxy- $C_{3-8}$ -cycloalk(en)yl, hydroxy- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, halo- $C_{1-6}$ -alk(en/yn)yl, halo- $C_{3-8}$ -cycloalk(en)yl, halo- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, cyano- $C_{1-6}$ -

alk(en/yn)yl, cyano-C<sub>3-8</sub>-cycloalk(en)yl and cyano-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, or


**R**<sup>12</sup> and **R**<sup>12'</sup> together with the nitrogen atom to which they are attached form a 4-8 membered saturated or unsaturated ring which optionally contains 1, 2 or 3

5 further heteroatoms;


with the proviso that when **R**<sup>3</sup> is **NR**<sup>12</sup>**R**<sup>12'</sup> then **q** is 0;

and


10 **Y** represents a group of formula **II**, **III**, **IV**, **V**, **VI**, **XXX** and **XXXI**:




II




III




IV



V




VI



XXX

or



XXXI

wherein

- 5 the line represents a bond attaching the group represented by Y to the carbon atom;

W is O or S;

- 10 T is N, NH or O;

L is N, C or CH;

a is 0, 1, 2 or 3;

5 b is 0, 1, 2, 3 or 4;

c is 0 or 1;

d is 0, 1, 2 or 3;

10

e is 0, 1 or 2;

f is 0, 1, 2, 3, 4 or 5;

15 g is 0, 1, 2, 3 or 4;

h is 0, 1, 2 or 3;

j is 0, 1, 2 or 3; with the proviso that when T is a nitrogen atom then j is 0, 1, 2 or  
20 3; and when T is NH or an oxygen atom then j is 0, 1 or 2;

k is 0, 1, 2, 3 or 4; and

each R<sup>5</sup> is independently selected from the group consisting of a C<sub>1-6</sub>-  
25 alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, Ar, Ar-  
C<sub>1-6</sub>-alk(en/yn)yl, Ar-thio, Ar-oxy, acyl, C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-  
cycloalk(en)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, halo-C<sub>1-6</sub>-  
alk(en/yn)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl,  
-CO-NR<sup>6</sup>R<sup>6'</sup>, cyano, cyano-C<sub>1-6</sub>-alk(en/yn)yl, cyano-C<sub>3-8</sub>-cycloalk(en)yl, cyano-  
30 C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, -NR<sup>7</sup>R<sup>7'</sup>, -S-R<sup>8</sup> and -SO<sub>2</sub>R<sup>8</sup>, or  
two adjacent R<sup>5</sup> together with the aromatic group to which they are attached form  
a 4-8 membered ring which optionally contains one or two heteroatoms;

$R^6$  and  $R^{6'}$  are independently selected from the group consisting of hydrogen,  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl and Ar;

5  $R^7$  and  $R^{7'}$  are independently selected from the group consisting of hydrogen,  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, Ar and acyl;

and

10  $R^8$  is selected from the group consisting of hydrogen,  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, Ar and  $-NR^9R^{9'}$ ; wherein  $R^9$  and  $R^{9'}$  are independently selected from the group consisting of hydrogen,  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl and  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl; provided that when  $R^8$  is  $-NR^9R^{9'}$  then  $R^5$  is not  $-S-R^8$ ;  
15 or salts thereof;

with the proviso that the compound of formula I is not:

N-[1-(phenylmethyl)-1H-indol-5-yl]-Methanesulfonamide;

N-[1-[(4-fluorophenyl)methyl]-1H-indol-5-yl]-Methanesulfonamide;

20 N-[2,3-dihydro-1-(phenylmethyl)-1H-indol-5-yl]-Methanesulfonamide;

N-[1-(phenylmethyl)-1H-indol-5-yl]-N'-4-quinolinyl-Urea;

N-[1-(phenylmethyl)-1H-indol-5-yl]-N'-4-quinolinyl-Urea; or

1-(1-benzyl-5-indoliny)-3-phenyl-Urea;

or salts thereof.

25

2. A compound according to Claim 1, wherein at least one of  $R^1$  or  $R^{1'}$  is a hydrogen atom.

30 3. A compound according to any one of Claims 1 and 2, wherein both  $R^1$  and  $R^{1'}$  are hydrogen atoms.

4. A compound according to any one of Claims 1-3, wherein s is 0.

5. A compound according to any one of Claims 1-3, wherein  $s$  is 1.
6. A compound according any one of Claims 1-5, wherein  $R^2$  is a hydrogen atom.
- 5 7. A compound according any one of Claims 1-4, wherein  $R^2$  is  $NO_2$  or a halogen atom.
8. A compound according to any one of Claims 1-3 and 5-7, wherein  $U$  is  $NR^{11}$ .
- 10 9. A compound according to Claim 8, wherein  $R^{11}$  is a hydrogen atom.
10. A compound according to any one of Claims 1-9, wherein  $X$  is  $CO$ .
11. A compound according to any one of Claims 1-9, wherein  $X$  is  $SO_2$ .
- 15 12. A compound according to any one of Claims 1-11, wherein  $q$  is 0.
13. A compound according to any one of Claims 1-11, wherein  $q$  is 1.
- 20 14. A compound according to Claim 13, wherein  $Z$  is an oxygen atom.
15. A compound according to any one of Claims 1-14, wherein  $R^3$  is selected from the group consisting of  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl, Ar, Ar- $C_{1-6}$ -alk(en/yn)yl, Ar-oxy- $C_{1-6}$ -alk(en/yn)yl, Ar- $C_{1-6}$ -alk(en/yn)yl oxy- $C_{1-6}$ -alk(en/yn)yl and  $-NR^{12}R^{12'}$ ; with the proviso that when  $R^3$  is  $NR^{12}R^{12'}$  then  $q$  is 0.
- 25 16. A compound according to Claim 15, wherein  $R^3$  is  $NR^{12}R^{12'}$ ,  $q$  is 0 and  $R^{12}$  and  $R^{12'}$  are independently selected from the group consisting of hydrogen,  $C_{1-6}$ -alk(en/yn)yl, Ar and Ar- $C_{1-6}$ -alk(en/yn)yl, or  $R^{12}$  and  $R^{12'}$  together with the  
30 nitrogen atom to which they are attached form a 4-8 membered saturated or unsaturated ring which optionally contains 1, 2 or 3 further heteroatoms.



17. A compound according to any one of Claims 1-16, wherein Y is of formula II, III, V, XXX, or XXXI.
18. A compound according to any of Claims 1-17, wherein Y is of formula II or III and W is a sulphur atom.
19. A compound according to any of Claims 1-17, wherein Y is of formula XXX and T is a nitrogen atom or an oxygen atom.
20. A compound according to any of Claims 1-17, wherein Y is of formula XXXI and L is C or CH.
21. A compound according to any of Claims 1-20, wherein each R<sup>5</sup> is independently selected from the group consisting of C<sub>1-6</sub>-alk(en/yn)yl, Ar, Ar-thio, Ar-oxy, halogen and halo-C<sub>1-6</sub>-alk(en/yn)yl or or two adjacent R<sup>5</sup> together with the aromatic group to which they are attached form a 4-8 membered ring which optionally contains one or two heteroatoms.
22. A compound according to any of Claims 1-21, said compounds being selected from the group consisting of:
- N*-[4-Chloro-1-(4-trifluoromethylbenzyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide;
- N*-[4-Chloro-1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide;
- [1-(4-Fluorobenzyl)-2,3-dihydro-1H-indol-5-yl]-carbamic acid propyl ester;
- N*-[1-(4-Fluorobenzyl)-2,3-dihydro-1H-indol-5-yl]-C-phenylmethanesulfonamide;
- 4-Fluoro-*N*-[1-(4-fluorobenzyl)-2,3-dihydro-1H-indol-5-yl]-benzamide;
- N*-[1-(4-Fluorobenzyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide;
- N*-[1-(4-Fluorobenzyl)-2,3-dihydro-1H-indol-5-yl]-2-thiophen-2-ylacetamide;
- N*-[1-(4-Fluorobenzyl)-2,3-dihydro-1H-indol-5-yl]-2-(4-fluorophenyl)-acetamide;
- 3-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-1,1-diisopropylurea;

- Morpholine-4-carboxylic acid [1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-amide;*
- Pyrrolidine-1-carboxylic acid [1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-amide;*
- 5     *[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-carbamic acid 2-benzyloxyethyl ester;*
- 3-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-1-methyl-1-propylurea;*
- [1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-carbamic acid*
- 10   *tert-butyl ester;*
- N-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-C-phenylmethanesulfonamide;*
- Butane-1-sulfonic acid [1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-amide;*
- 15   *N-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-4-fluorobenzamide;*
- N-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-2,2-dimethylpropionamide;*
- N-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-2-*
- 20   *phenoxyacetamide;*
- N-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide;*
- N-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-butyramide;*
- Cyclopentanecarboxylic acid [1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-*
- 25   *indol-5-yl]-amide;*
- N-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-2-thiophen-2-ylacetamide;*
- N-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-isonicotinamide;*
- 30   *N-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-4-dimethylaminobenzamide;*
- N-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-2-(4-fluorophenyl)-acetamide;*


- N*-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-6-trifluoromethylnicotinamide;
- 1-*tert*-Butyl-3-[1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-urea;
- 5    1-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-3-ethylurea;
- 1-Benzyl-3-[1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-urea;
- 1-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-3-phenethylurea;
- 1-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-3-thiophen-2-ylurea;
- 10    1-[1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-3-thiophen-3-ylurea;
- [1-(5-Chlorothiophen-2-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-carbamic acid propyl ester;
- 15    2,2-Dimethyl-*N*-[6-nitro-1-(4-trifluoromethylbenzyl)-2,3-dihydro-1*H*-indol-5-yl]-propionamide;
- N*-[1-(5-Chlorothiophen-2-ylmethyl)-6-nitro-2,3-dihydro-1*H*-indol-5-yl]-2,2-dimethylpropionamide;
- 2-(4-Fluorophenyl)-*N*-[6-nitro-1-(4-trifluoromethylbenzyl)-2,3-dihydro-1*H*-indol-5-yl]-acetamide;
- 20    5-yl]-acetamide;
- N*-[1-(5-Chlorothiophen-2-ylmethyl)-6-nitro-2,3-dihydro-1*H*-indol-5-yl]-2-(4-fluorophenyl)-acetamide;
- N*-[1-(5-Chlorothiophen-2-ylmethyl)-6-nitro-2,3-dihydro-1*H*-indol-5-yl]-3,3-dimethylbutyramide;
- 25    *N*-[6-Amino-1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-3,3-dimethylbutyramide;
- N*-[6-Amino-1-(4-trifluoromethylbenzyl)-2,3-dihydro-1*H*-indol-5-yl]-2,2-dimethylpropionamide;
- N*-[6-Amino-1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-2,2-dimethylpropionamide;
- 30    2-(4-fluorophenyl)-acetamide;
- N*-[6-Amino-1-(4-trifluoromethylbenzyl)-2,3-dihydro-1*H*-indol-5-yl]-2-(4-fluorophenyl)-acetamide;

- N*-[6-Amino-1-(4-trifluoromethylbenzyl)-2,3-dihydro-1*H*-indol-5-yl]-3,3-dimethylbutyramide;
- N*-[6-Amino-1-(4-fluorobenzyl)-2,3-dihydro-1*H*-indol-5-yl]-3,3-dimethylbutyramide;
- 5 *N*-[6-Amino-1-(3-fluoro-4-trifluoromethylbenzyl)-2,3-dihydro-1*H*-indol-5-yl]-3,3-dimethylbutyramide;
- N*-[1-(5-Chlorothiophen-2-ylmethyl)-1*H*-indol-5-yl]-3,3-dimethylbutyramide;
- N*-[6-Bromo-1-(4-trifluoromethylbenzyl)-2,3-dihydro-1*H*-indol-5-yl]-3,3-dimethylbutyramide;
- 10 *N*-[6-Bromo-1-(5-chlorothiophen-2-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-3,3-dimethylbutyramide;
- N*-[1-(4-Chlorobenzyl)-2,3-dihydro-1*H*-indol-5-yl]-3,3-dimethylbutyramide;
- 3,3-Dimethyl-*N*-[1-(4-trifluoromethylbenzyl)-2,3-dihydro-1*H*-indol-5-yl]-butyramide;
- 15 *N*-[1-(4-Isopropylbenzyl)-2,3-dihydro-1*H*-indol-5-yl]-3,3-dimethylbutyramide;
- N*-[1-(3-Fluoro-4-trifluoromethylbenzyl)-2,3-dihydro-1*H*-indol-5-yl]-3,3-dimethylbutyramide;
- N*-[1-(6-Chlorobenzo[1,3]dioxol-5-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-3,3-dimethylbutyramide;
- 20 *N*-[1-(3,5-Dimethyl-1-phenyl-1*H*-pyrazol-4-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-3,3-dimethylbutyramide;
- N*-[1-(2-Chloro-5-trifluoromethylbenzyl)-2,3-dihydro-1*H*-indol-5-yl]-3,3-dimethylbutyramide;
- N*-{1-[5-(4-Chlorophenoxy)-1,3-dimethyl-1*H*-pyrazol-4-ylmethyl]-2,3-dihydro-1*H*-indol-5-yl}-3,3-dimethylbutyramide;
- 25 3,3-Dimethyl-*N*-[1-(6-*p*-tolylloxy-pyridin-3-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-butyramide;
- N*-{1-[6-(4-Chlorophenylsulfanyl)-pyridin-3-ylmethyl]-2,3-dihydro-1*H*-indol-5-yl}-3,3-dimethylbutyramide;
- 30 *N*-{1-[6-(4-Cyanophenoxy)-pyridin-3-ylmethyl]-2,3-dihydro-1*H*-indol-5-yl}-3,3-dimethylbutyramide;
- 3,3-Dimethyl-*N*-[1-(6-trifluoromethylpyridin-3-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-butyramide;

- 3,3-Dimethyl-N-[1-(3-methyl-benzo[b]thiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-butyramide;*
- N-[1-(6-Fluoro-4H-benzo[1,3]dioxin-8-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-3,3-dimethylbutyramide;*
- 5 *3,3-Dimethyl-N-[1-(6-phenoxy-pyridin-3-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-butyramide;*
- 3,3-Dimethyl-N-[1-(3-methyl-5-phenyl-isoxazol-4-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-butyramide;*
- N-(1-Benzo[b]thiophen-2-ylmethyl-2,3-dihydro-1H-indol-5-yl)-3,3-dimethylbutyramide;*
- 10 *N-{1-[1-(4-Fluorophenyl)-5-methyl-1H-pyrazol-4-ylmethyl]-2,3-dihydro-1H-indol-5-yl}-3,3-dimethylbutyramide;*
- 3,3-Dimethyl-N-[1-(5-methylthiophen-2-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-butyramide;*
- 15 *3,3-Dimethyl-N-[1-(4-pyrrol-1-yl-benzyl)-2,3-dihydro-1H-indol-5-yl]-butyramide;*
- N-[1-(4-Chlorobenzyl)-2,3-dihydro-1H-indol-5-yl]-2-(4-fluorophenyl)-acetamide;*
- 2-(4-Fluorophenyl)-N-[1-(4-trifluoromethylbenzyl)-2,3-dihydro-1H-indol-5-yl]-acetamide;*
- 20 *2-(4-Fluorophenyl)-N-[1-(4-isopropylbenzyl)-2,3-dihydro-1H-indol-5-yl]-acetamide;*
- 2-(4-Fluorophenyl)-N-[1-(3-fluoro-4-trifluoromethylbenzyl)-2,3-dihydro-1H-indol-5-yl]-acetamide;*
- 25 *N-[1-(6-Chlorobenzo[1,3]dioxol-5-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-2-(4-fluorophenyl)-acetamide;*
- N-[1-(3,5-Dimethyl-1-phenyl-1H-pyrazol-4-ylmethyl)-2,3-dihydro-1H-indol-5-yl]-2-(4-fluorophenyl)-acetamide;*
- N-[1-(2-Chloro-5-trifluoromethylbenzyl)-2,3-dihydro-1H-indol-5-yl]-2-(4-fluorophenyl)-acetamide;*
- 30 *N-{1-[5-(4-Chlorophenoxy)-1,3-dimethyl-1H-pyrazol-4-ylmethyl]-2,3-dihydro-1H-indol-5-yl}-2-(4-fluorophenyl)-acetamide;*

- N*-{1-[6-(4-Cyanophenoxy)-pyridin-3-ylmethyl]-2,3-dihydro-1*H*-indol-5-yl}-2-(4-fluorophenyl)-acetamide;
- 2-(4-Fluorophenyl)-*N*-[1-(3-methyl-benzo[*b*]thiophen-2-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-acetamide;
- 5 *N*-[1-(6-Fluoro-4*H*-benzo[1,3]dioxin-8-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-2-(4-fluorophenyl)-acetamide;
- 2-(4-Fluorophenyl)-*N*-[1-(6-phenoxy-pyridin-3-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-acetamide;
- 10 *N*-(1-Benzo[*b*]thiophen-2-ylmethyl-2,3-dihydro-1*H*-indol-5-yl)-2-(4-fluorophenyl)-acetamide;
- 2-(4-Fluorophenyl)-*N*-{1-[1-(4-fluorophenyl)-5-methyl-1*H*-pyrazol-4-ylmethyl]-2,3-dihydro-1*H*-indol-5-yl}-acetamide;
- 2-(4-Fluorophenyl)-*N*-[1-(5-methylthiophen-2-ylmethyl)-2,3-dihydro-1*H*-indol-5-yl]-acetamide; and
- 15 2-(4-Fluorophenyl)-*N*-[1-(4-pyrrol-1-yl-benzyl)-2,3-dihydro-1*H*-indol-5-yl]-acetamide, or
- a pharmaceutically acceptable salt thereof.

23. A pharmaceutical composition comprising one or more pharmaceutically acceptable carriers or diluents and a compound according to any one of claims 1-22.
- 20
24. Use of a pharmaceutical composition comprising one or more pharmaceutically acceptable carriers or diluents and a compound of the general formula I
- 25



(I)

wherein

the dotted line represents an optional bond;

5

$R^1$  and  $R^{1'}$  are independently selected from the group consisting of hydrogen,  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, hydroxy- $C_{1-6}$ -alk(en/yn)yl, hydroxy- $C_{3-8}$ -cycloalk(en)yl, hydroxy- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, halo- $C_{1-6}$ -alk(en/yn)yl, halo- $C_{3-8}$ -cycloalk(en)yl, halo- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, cyano- $C_{1-6}$ -alk(en/yn)yl, cyano- $C_{3-8}$ -cycloalk(en)yl and cyano- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl; or  $R^1$  and  $R^{1'}$  together with the carbon atom to which they are attached form a 3-8 membered saturated or unsaturated ring which optionally contains 1 or 2 heteroatoms;

15

s is 0 or 1;

U is O,  $NR^{11}$ , S,  $SO_2$ ,  $SO_2NR^{11}$ , CO-O or CO- $NR^{11}$ ; wherein  $R^{11}$  is selected from the group consisting of hydrogen,  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl; or  $R^2$  and  $R^{11}$  together with the nitrogen atom to which they are attached form a 4-8 membered saturated or unsaturated ring which optionally contains 1, 2 or 3 further heteroatoms;

20

$R^2$  is selected from the group consisting of hydrogen,  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, Ar, Ar- $C_{1-6}$ -alk(en/yn)yl, Ar- $C_{3-8}$ -cycloalk(en)yl, Ar- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, acyl, hydroxy- $C_{1-6}$ -alk(en/yn)yl, hydroxy- $C_{3-8}$ -cycloalk(en)yl, hydroxy- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, halogen, halo- $C_{1-6}$ -alk(en/yn)yl, halo- $C_{3-8}$ -cycloalk(en)yl, halo- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, cyano, cyano- $C_{1-6}$ -alk(en/yn)yl, cyano- $C_{3-8}$ -cycloalk(en)yl, cyano- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl,  $-NO_2$ ,  $NR^{10}R^{10'}$ - $C_{1-6}$ -alk(en/yn)yl,  $NR^{10}R^{10'}$ - $C_{3-8}$ -cycloalk(en)yl and  $NR^{10}R^{10'}$ - $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl; wherein

25

30

$R^{10}$  and  $R^{10'}$  are independently selected from the group consisting of hydrogen,  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, hydroxy- $C_{1-6}$ -alk(en/yn)yl, hydroxy- $C_{3-8}$ -cycloalk(en)yl, hydroxy- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, halo- $C_{1-6}$ -alk(en/yn)yl, halo- $C_{3-8}$ -cycloalk(en)yl, halo- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, cyano- $C_{1-6}$ -alk(en/yn)yl, cyano- $C_{3-8}$ -cycloalk(en)yl and cyano- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, or

$R^{10}$  and  $R^{10'}$  together with the nitrogen atom to which they are attached form a 4-8 membered saturated or unsaturated ring which optionally contains 1, 2 or 3 further heteroatoms;

with the proviso that when  $R^2$  is  $NO_2$ , halogen or cyano then s is 0; and with the proviso that when  $R^2$  is a hydrogen atom or acyl and s is 1 then U is  $NR^{11}$ , O or S;

wherein the group  $-(U)_s-R^2$  is linked to position 4 or 6 of the indole or indoline;

q is 0 or 1;

Z is O or S;

X is CO or  $SO_2$ ; with the proviso that q is 0 when X is  $SO_2$ ;







$R^3$  is selected from the group consisting of  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl, heterocycloalk(en)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl,  $C_{1-6}$ -alk(en/yn)yl- $C_{3-8}$ -cycloalk(en)yl,  $C_{1-6}$ -alk(en/yn)yl-heterocycloalk(en)yl, Ar, Ar- $C_{1-6}$ -alk(en/yn)yl, Ar- $C_{3-8}$ -cycloalk(en)yl, Ar-heterocycloalk(en)yl, Ar- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, Ar- $C_{1-6}$ -alk(en/yn)yl- $C_{3-8}$ -cycloalk(en)yl, Ar- $C_{1-6}$ -alk(en/yn)yl-heterocycloalk(en)yl,  $C_{1-6}$ -alk(en/yn)yl-oxy- $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl-oxy- $C_{1-6}$ -alk(en/yn)yl,  $C_{1-6}$ -alk(en/yn)yl-oxy- $C_{3-8}$ -cycloalk(en)yl,  $C_{1-6}$ -alk(en/yn)yl-oxy-heterocycloalk(en)yl, Ar-oxy- $C_{1-6}$ -alk(en/yn)yl, Ar- $C_{1-6}$ -alk(en/yn)yl-oxy- $C_{1-6}$ -alk(en/yn)yl,  $C_{1-6}$ -alk(en/yn)yl-oxy-carbonyl- $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl-oxy-carbonyl- $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl-oxy-carbonyl- $C_{1-6}$ -alk(en/yn)yl, hydroxy- $C_{1-6}$ -alk(en/yn)yl, hydroxy- $C_{3-8}$ -cycloalk(en)yl, hydroxy-heterocycloalk(en)yl,




hydroxy-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, hydroxy-C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl, hydroxy-C<sub>1-6</sub>-alk(en/yn)yl-heterocycloalk(en)yl, halo-C<sub>1-6</sub>-alk(en/yn)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl, halo-heterocycloalk(en)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, halo-C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl, halo-C<sub>1-6</sub>-alk(en/yn)yl-heterocycloalk(en)yl, halo-C<sub>1-6</sub>-alk(en/yn)yl-Ar, halo-C<sub>3-8</sub>-cycloalk(en)yl-Ar, halo-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl-Ar, halo-C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl-Ar, cyano-C<sub>1-6</sub>-alk(en/yn)yl, cyano-C<sub>3-8</sub>-cycloalk(en)yl, cyano-heterocycloalk(en)yl, cyano-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, cyano-C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl, cyano-C<sub>1-6</sub>-alk(en/yn)yl-heterocycloalk(en)yl, acyl-C<sub>1-6</sub>-alk(en/yn)yl, acyl-C<sub>3-8</sub>-cycloalk(en)yl, acyl-heterocycloalk(en)yl, acyl-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, acyl-C<sub>1-6</sub>-alk(en/yn)yl-C<sub>3-8</sub>-cycloalk(en)yl, acyl-C<sub>1-6</sub>-alk(en/yn)yl-heterocycloalk(en)yl and -NR<sup>12</sup>R<sup>12'</sup>, optionally substituted NR<sup>12</sup>R<sup>12'</sup>-C<sub>1-6</sub>-alk(en/yn)yl, optionally substituted NR<sup>12</sup>R<sup>12'</sup>-C<sub>3-8</sub>-cycloalk(en)yl, optionally substituted NR<sup>12</sup>R<sup>12'</sup>-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl; wherein R<sup>12</sup> and R<sup>12'</sup> are independently selected from the group consisting of hydrogen, C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, Ar, Ar-C<sub>1-6</sub>-alk(en/yn)yl, Ar-C<sub>3-8</sub>-cycloalk(en)yl, Ar-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, hydroxy-C<sub>1-6</sub>-alk(en/yn)yl, hydroxy-C<sub>3-8</sub>-cycloalk(en)yl, hydroxy-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, halo-C<sub>1-6</sub>-alk(en/yn)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl, halo-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, cyano-C<sub>1-6</sub>-alk(en/yn)yl, cyano-C<sub>3-8</sub>-cycloalk(en)yl and cyano-C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, or R<sup>12</sup> and R<sup>12'</sup> together with the nitrogen atom to which they are attached form a 4-8 membered saturated or unsaturated ring which optionally contains 1, 2 or 3 further heteroatoms; with the proviso that when R<sup>3</sup> is NR<sup>12</sup>R<sup>12'</sup> then q is 0;

and

Y represents a group of formula II, III, IV, V, , VI, XXX and XXXI:

**II****III****IV****V****VI****XXX**

or

**XXXI**

wherein

- 5 the line represents a bond attaching the group represented by Y to the carbon atom;

W is O or S;

- 10 T is N, NH or O;

L is N, C or CH;

a is 0, 1, 2 or 3;

5 b is 0, 1, 2, 3 or 4;

c is 0 or 1;

d is 0, 1, 2 or 3;

10

e is 0, 1 or 2;

f is 0, 1, 2, 3, 4 or 5;

15

g is 0, 1, 2, 3 or 4;

h is 0, 1, 2 or 3;

20

j is 0, 1, 2 or 3; with the proviso that when T is a nitrogen atom then j is 0, 1, 2 or 3; and when T is NH or an oxygen atom then j is 0, 1 or 2;

k is 0, 1, 2, 3 or 4; and

25

each  $R^5$  is independently selected from the group consisting of a  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, Ar, Ar- $C_{1-6}$ -alk(en/yn)yl, Ar-thio, Ar-oxy, acyl,  $C_{1-6}$ -alk(en/yn)yl-oxy,  $C_{3-8}$ -cycloalk(en)yl-oxy,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl-oxy, halogen, halo- $C_{1-6}$ -alk(en/yn)yl, halo- $C_{3-8}$ -cycloalk(en)yl, halo- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, -CO-NR<sup>6</sup>R<sup>6'</sup>, cyano, cyano- $C_{1-6}$ -alk(en/yn)yl, cyano- $C_{3-8}$ -cycloalk(en)yl, cyano- $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, -NR<sup>7</sup>R<sup>7'</sup>, -S-R<sup>8</sup> and -SO<sub>2</sub>R<sup>8</sup>, or  
30 two adjacent  $R^5$  together with the aromatic group to which they are attached form a 4-8 membered ring which optionally contains one or two heteroatoms;

$R^6$  and  $R^{6'}$  are independently selected from the group consisting of hydrogen,  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl and Ar;

5  $R^7$  and  $R^{7'}$  are independently selected from the group consisting of hydrogen,  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, Ar and acyl;

and

10  $R^8$  is selected from the group consisting of hydrogen,  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, Ar and  $-NR^9R^{9'}$ ; wherein  $R^9$  and  $R^{9'}$  are independently selected from the group consisting of hydrogen,  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl and  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl; provided that when  $R^8$  is  $-NR^9R^{9'}$  then  $R^5$  is not  $-S-R^8$ ;

15

or salts thereof

for increasing ion flow in a potassium channel of a mammal such as a human.

20 25. Use according to Claim 24 for the prevention, treatment or inhibition of a disorder or condition being responsive to an increased ion flow in a potassium channel, such disorder or condition is preferably a disorder or condition of the central nervous system.

25 26. Use according to Claim 25, wherein said disorder or disease is selected from the group consisting of seizure disorders such as convulsions, epilepsy and status epilepticus.

30 27. Use according to Claim 25, characterized in that the disorder or condition is selected from the group consisting of neuropathic and migraine pain disorders such as allodynia, hyperalgesic pain, phantom pain, neuropathic pain related to diabetic neuropathy and neuropathic pain related to migraine

28. Use according to Claim 25, characterized in that the disorder or condition is selected from the group consisting of anxiety disorders such as anxiety, generalized anxiety disorder, panic anxiety, obsessive compulsive disorder, social phobia, performance anxiety, post-traumatic stress disorder, acute stress reaction, adjustment disorders, hypochondriacal disorders, separation anxiety disorder, agoraphobia, specific phobias, anxiety disorder due to general medical condition and substance-induced anxiety disorder.
29. Use according to Claim 25, characterized in that the disorder or condition is selected from the group consisting of and neurodegenerative disorders such as Alzheimer's disease, Huntington's chorea, multiple sclerosis, amyotrophic lateral sclerosis, AIDS-induced encephalopathy and other infection-related encephalopathies being caused by rubella viruses, herpes viruses, borrelia and by unknown pathogens, Creutzfeld-Jakob disease, Parkinson's disease, trauma-induced neurodegenerations.
30. Use according to Claim 25, characterized in that the disorder or condition is selected from the group consisting of neuronal hyperexcitation states such as in medicament withdrawal or by intoxication.

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D209/08 C07D409/06 C07D401/06 C07D403/06 C07D407/06  
C07D417/06 A61K31/404 A61P25/08

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BEILSTEIN Data, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with Indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 03/031409 A (CHEIL JEDANG CORP) 17 April 2003 (2003-04-17) page 7; examples ---	1-30
X	WO 01/12187 A (HARGREAVES RODNEY BRIAN ; WHITTAMORE PAUL ROBERT OWEN (GB); ASTRAZENECA) 22 February 2001 (2001-02-22) examples 25, 33, 51, 55, 85 ---	1-16, 21, 23-30
A	WO 99/09024 A (JOHNS AMANDA ; PORTER RODERICK ALAN (GB); SMITHKLINE BEECHAM PLC (GB);) 25 February 1999 (1999-02-25) page 27; examples ---	1-30
A	US 5 234 942 A (BERNSTEIN PETER R ET AL) 10 August 1993 (1993-08-10) column 56 -column 59; examples --- -/--	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*&amp;\* document member of the same patent family

Date of the actual completion of the international search

23 July 2004

Date of mailing of the international search report

24/08/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Menegaki, F

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	RODERICK A. ET AL: "1,3-Biarylureas as selective non-peptide antagonists of the Orexin-1 receptor" BIOORGANIC MEDICINAL CHEMISTRY LETTERS, vol. 11, no. 14, 23 July 2001 (2001-07-23), pages 1907-1910, XP002289307 example 3; table 1 ---	1-30
A	WO 02/066426 A (SQUIBB BRISTOL MYERS CO) 29 August 2002 (2002-08-29) examples ---	1-30
P,X	LACONDE, G. ET AL: "New Analogues of Anticancer E7070" JOURNAL OF ENZYME INHIBITION AND MEDICINAL CHEMISTRY, vol. 18, no. 2, 2003, pages 89-94, XP009034199 page 93; figure 1; example 7 -----	1-16,21, 23-30

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 03031409	A	17-04-2003	KR 2003030382 A EP 1442016 A1 WO 03031409 A1 US 2003109568 A1	18-04-2003 04-08-2004 17-04-2003 12-06-2003
WO 0112187	A	22-02-2001	AU 6583400 A BR 0013368 A CA 2380775 A1 CN 1379774 T EP 1210343 A2 WO 0112187 A2 JP 2003507327 T NO 20020765 A NZ 517059 A ZA 200200669 A	13-03-2001 07-05-2002 22-02-2001 13-11-2002 05-06-2002 22-02-2001 25-02-2003 17-04-2002 28-05-2004 24-04-2003
WO 9909024	A	25-02-1999	AU 8741198 A CA 2300178 A1 EP 1003737 A1 WO 9909024 A1 JP 2001515075 T US 6410529 B1	08-03-1999 25-02-1999 31-05-2000 25-02-1999 18-09-2001 25-06-2002
US 5234942	A	10-08-1993	AT 56205 T AU 583062 B2 AU 4881485 A CA 1273934 A1 CN 85108623 A DD 253618 A5 DE 3579554 D1 DK 479385 A EP 0179619 A1 ES 8704458 A1 ES 8800899 A1 ES 8800900 A1 ES 8802495 A1 ES 8801786 A1 FI 854024 A GR 852519 A1 HU 38905 A2 HU 194163 B IE 58848 B1 IL 76756 A JP 2015992 C JP 7045466 B JP 61178963 A KR 9007419 B1 MW 3285 A1 NO 854163 A NZ 213872 A PL 255841 A1 PT 81297 A, B SU 1545940 A3 US 4997844 A ZA 8507952 A ZM 7885 A1 ZW 18185 A1 PL 265549 A1	15-09-1990 20-04-1989 24-04-1986 11-09-1990 30-07-1986 27-01-1988 11-10-1990 20-04-1986 30-04-1986 16-06-1987 16-02-1988 16-02-1988 16-10-1988 01-05-1988 20-04-1986 24-01-1986 28-07-1986 28-01-1988 17-11-1993 15-05-1989 19-02-1996 17-05-1995 11-08-1986 08-10-1990 19-06-1987 21-04-1986 28-11-1989 21-09-1987 01-11-1985 23-02-1990 05-03-1991 28-05-1986 21-02-1986 20-05-1987 01-09-1988



Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5234942	A	SU 1595338 A3	23-09-1990
WO 02066426	A	29-08-2002	
		CA 2438805 A1	29-08-2002
		EP 1363624 A2	26-11-2003
		HU 0400063 A2	28-04-2004
		WO 02066426 A2	29-08-2002
		US 2002156120 A1	24-10-2002